

Harmonic analysis on local fields and adelic spaces I

D.V. Osipov, A.N. Parshin *

Abstract

Harmonic analysis is developed on objects of some category C_2 of infinite-dimensional filtered vector spaces over a finite field. This category includes two-dimensional local fields and adelic spaces of algebraic surfaces defined over a finite field. The main result is the theory of the Fourier transform on these objects and obtaining of two-dimensional Poisson formulas.

1 Introduction

Local fields and adelic groups composed from them are well known tools of arithmetic. They were introduced by C. Chevalley in the 1930s and were used to formulate and solve many different problems in number theory and algebraic geometry (see, for example, [1, 20]). These constructions are associated with fields of algebraic numbers and fields of algebraic functions of one variable, i.e. with schemes of dimension 1. A need for such constructions in higher dimensions was realized by A. N. Parshin in the 1970s. He managed to develop these constructions in the local case for any dimension and in the global case for dimension 2, [14]. For the schemes of an arbitrary dimension this approach was developed by A. A. Beilinson, [2].

If X is a scheme of dimension n and of finite type over \mathbb{Z} , and $X_n \subset X_{n-1} \subset \dots \subset X_1 \subset X_0 \subset X$ is a flag of irreducible subschemes ($\text{codim}(X_i) = i$), then one can define a ring $K_{X_0, \dots, X_{n-1}}$ associated to the flag. In case when all the subschemes are regularly embedded, the ring is an n -dimensional local field. Then one can form an adelic object

$$\mathbb{A}_X = \prod' K_{X_0, \dots, X_{n-1}}$$

where the product is taken over all the flags, but with certain restrictions on the components of adeles (see [2, 6, 14, 4, 12]). For a scheme over a finite field \mathbb{F}_q this definition is the ultimate definition of the adelic space attached to X . In general, one has to extend it to $\mathbb{A}_X \oplus \mathbb{A}_{X \otimes \mathbb{R}}$ by adding archimedean components.

In dimension 1, the adelic groups \mathbb{A}_X and \mathbb{A}_X^* are locally compact groups and thus we can develop the classical harmonic analysis on these groups. The starting

*Both authors are supported by RFBR (grant no. 08-01-00095-a), by a program of President of RF for supporting of Leading Scientific Schools (grant no. 1987.2008.1), by INTAS (grant no. 05-1000008-8118); besides the first author is supported by a grant of Russian Science Support Foundation and by a program of President of RF for supporting of young russian scientists (grant no. MK-864.2008.1).

point for that is the measure theory on locally compact local fields attached to points on schemes X of dimension 1.

This harmonic analysis can be applied to a study of zeta- and L -functions of one-dimensional schemes X . J. Tate and independently K. Iwasawa have introduced an analytically defined L -function

$$L(s, \chi, f) = \int_{\mathbb{A}^*} \chi(a) |a|^s f(a) d^* \mu,$$

where $d^* \mu$ is a Haar measure on \mathbb{A}_X^* , the function f belongs to the Bruhat-Schwartz space of functions on \mathbb{A}_X and χ is an abelian character, coming from the Galois group $\text{Gal}(K^{\text{ab}}/K)$ by the reciprocity map

$$\mathbb{A}_X^* \rightarrow \text{Gal}(K^{\text{ab}}/K).$$

For $L(s, \chi, f)$, they have proved an analytical continuation to the whole s -plane and the functional equation

$$L(s, \chi, f) = L(1-s, \chi^{-1}, \hat{f}),$$

using Fourier transform $f \mapsto \hat{f}$ and Poisson formula for functions f on the space \mathbb{A}_X [1, 20]. If $L_X(s, \chi)$ is the arithmetically defined L -function of the scheme X , then $L_X(s, \chi) = L(s, \chi, f)$ for an appropriate function f .

For a long time, the second author had been pointing to an essential gap in arithmetics related with absence of such approach for schemes of dimension > 1 and suggesting that the gap should be filled. We note that there is a cohomological formalism defined for schemes over \mathbb{F}_q and working equally well in all dimensions.

Problem 1. *Extend Tate–Iwasawa’s analytic method to higher dimensions* (see, in particular, [17, 18]).

The higher adeles have been introduced exactly for this purpose. So we have the following underlying

Problem 2. *Develop a measure theory and harmonic analysis on n -dimensional local fields.*

Note that the n -dimensional local fields are not locally compact topological spaces for $n > 1$ and by Weil’s theorem the existence of the Haar measure (in the usual sense) on a topological group implies its locally compactness.

In the paper, we show how to construct a harmonic analysis and a measure theory in the first non-trivial case of algebraic surface X over a finite field k (or two-dimensional schemes over \mathbb{Z}). In this situation, the corresponding adelic objects will look as follows. Let P be a closed point of X , $C \subset X$ be an irreducible curve such that $P \in C$.

If X and C are smooth at P then let $t \in \mathcal{O}_{X,P}$ be a local equation of C at P and $u \in \mathcal{O}_{X,P}$ be such that $u|_C \in \mathcal{O}_{C,P}$ is a local parameter at P . Denote by \wp the ideal defining the curve C near P . Now we can introduce the two-dimensional local field attached to the pair P, C by the procedure including completions and localizations: $K_{P,C} = \text{Frac}((\widehat{\mathcal{O}_{X,P}})_{\wp}) = \text{Frac}(\widehat{\mathcal{O}_{x,C}}) = k(P)((u))((t))$. This definition can be easily extended to the case of a singular point P on the curve C . Then one can take the adelic product of the local fields over all the pairs P, C . We get the adelic space \mathbb{A}_X on the surface X .

If C is a curve then the space \mathbb{A}_C contains important subspaces $\mathbb{A}_0 = K = k(C)$ of principal adeles and $\mathbb{A}_1 = \prod_{x \in C} \hat{\mathcal{O}}_x$ of integral adeles, from which one constructs the adelic complex: $\mathbb{A}_0 \oplus \mathbb{A}_1 \rightarrow \mathbb{A}_C$. In dimension 2, there is a much more complicated structure of subspaces in \mathbb{A}_X . Among the others, it includes subspaces $\mathbb{A}_{12} = \prod'_{x \in C} \hat{\mathcal{O}}_{x,C}$ and, more generally, $\mathbb{A}_{12}(D)$, where the divisor D indicates on some regularity condition around the D (see [4] and a brief exposition in [15, 18]).

In dimension 1, the group \mathbb{A}_C is a locally compact group and the spaces \mathbb{A}_0 and \mathbb{A}_1 are, correspondingly, discrete and compact. The construction of analysis starts with a definition of functional spaces. Let V be a finite dimensional vector space either over an adelic ring \mathbb{A}_C , or over an one-dimensional local field K with finite residue field \mathbb{F}_q , or a subspace of type \mathbb{A}_0 or \mathbb{A}_1 . We put

$$\begin{aligned} \mathcal{D}(V) &= \{\text{locally constant functions with compact support}\} \\ \tilde{\mathcal{E}}(V) &= \{\text{uniformly locally constant functions}\} \\ \mathcal{E}(V) &= \{\text{all locally constant functions}\} \\ \mathcal{D}'(V) &= \{\text{dual to } \mathcal{D}, \text{ i.e. all distributions}\} \\ \tilde{\mathcal{E}}'(V) &= \{\text{"continuously" dual to } \tilde{\mathcal{E}}\} \\ \mathcal{E}'(V) &= \{\text{"continuosulsy" dual to } \mathcal{E}, \text{ i.e. distributions with compact support}\}. \end{aligned}$$

These are the classical spaces introduced by F. Bruhat [3]. The harmonic analysis includes definitions of direct and inverse images in some category C_1 of the spaces like V , a definition of the Fourier transform \mathbf{F} as a map from $\mathcal{D}'(V) \otimes \mu(V)^*$ to $\mathcal{D}'(\check{V})$ as well for the other types of spaces. Here $\mu(V)$ is a space of the Haar measures on V and \check{V} is a dual object (continuous linear functionals from the full dual space). The main result is the following Poisson formula

$$\mathbf{F}(\delta_{W, \mu_0} \otimes \mu^{-1}) = \delta_{W^\perp \cap \check{V}, \mu^{-1}/\mu_0^{-1}}$$

for any closed subgroup $i : W \rightarrow V$. Here $\mu_0 \in \mu(W)$, $\mu \in \mu(V)$, $\delta_{W, \mu_0} = i_*(\mu_0)$ (we use that, by definition of measures, $\mu(W) \subset \mathcal{D}'(W)$, and the direct image i_* maps $\mathcal{D}'(W)$ to $\mathcal{D}'(V)$), and $W^\perp \cap \check{V}$ is the annihilator of W in \check{V} .

All these constructions are carefully done in section 4 (in section 3, we collected the corresponding facts for the case of dimension 0, section 2 contains the basic notations used throughout the paper). The main issues here are the theory of C_0 and C_1 spaces (section 4.1), definition of the spaces of functions and distributions \mathcal{D} , \mathcal{D}' , \mathcal{E} , \mathcal{E}' , $\tilde{\mathcal{E}}$, $\tilde{\mathcal{E}}'$ (section 4.2), invariant Haar measures (section 4.3), Fourier transform (section 4.4), Poisson formula (section 4.5), Fubini formula and table of all possible direct and inverse images for the morphisms in the category C_1 (section 4.6), and the base change rules (section 4.7). The latter one is very useful when we extend the theory to the higher dimensions. The exposition is completely independent of the standard ones.

This general formalism can be applied to the adelic space on a curve C and we immediately get the following facts of analysis on the self-dual group \mathbb{A}_C : $\mathbf{F}(\delta_{\mathbb{A}_1(D)}) = \text{vol}(\mathbb{A}_1(D))\delta_{\mathbb{A}_1((\omega)-D)}$, and $\mathbf{F}(\delta_K) = \text{vol}(\mathbb{A}/K)^{-1}\delta_K$ for the standard subgroups in \mathbb{A}_C (attached to a divisor D and to the principal adeles, respectively). This easily implies Riemann-Roch and Serre duality for divisors on the curve C (see[18]).

For dimension 2, the space \mathbb{A}_X has a filtration by the subspaces $\mathbb{A}_{12}(D)$ where D runs through the Cartier divisors on X . The quotients $\mathbb{A}_{12}(D)/\mathbb{A}_{12}(D')$ will be spaces of the type considered above in the one-dimensional situation. This allows us to introduce the following spaces of functions (distributions):

$$\begin{aligned}
\mathcal{D}_{P_0}(V) &= \lim_{\leftarrow j^*} \lim_{\leftarrow i_*} \mathcal{D}(P/Q) \otimes \mu(P_0/Q), \\
\mathcal{D}'_{P_0}(V) &= \lim_{\rightarrow j_*} \lim_{\rightarrow i^*} \mathcal{D}'(P/Q) \otimes \mu(P_0/Q)^*, \\
\mathcal{E}(V) &= \lim_{\leftarrow j^*} \lim_{\rightarrow i^*} \mathcal{E}(P/Q), \\
\mathcal{E}'(V) &= \lim_{\rightarrow j_*} \lim_{\leftarrow i_*} \mathcal{E}'(P/Q), \\
\tilde{\mathcal{E}}(V) &= \lim_{\rightarrow i^*} \lim_{\leftarrow j^*} \tilde{\mathcal{E}}(P/Q), \\
\tilde{\mathcal{E}}'(V) &= \lim_{\leftarrow i_*} \lim_{\rightarrow j_*} \tilde{\mathcal{E}}'(P/Q),
\end{aligned}$$

where $P \supset Q \supset R$ are some elements of the filtration in \mathbb{A}_X (more generally, in a reasonable filtered space V with locally compact quotients), P_0 is a fixed subspace from the filtration and $j : Q/R \rightarrow P/R$, $i : P/R \rightarrow P/Q$ are the canonical maps. Note that both spaces $\mathcal{D}_{P_0}(V)$, $\mathcal{D}'_{P_0}(V)$ are $\mathcal{E}(V)$ -modules. In contrast to the one-dimensional case, there are no simple inclusions between these spaces.

Just as in the case of dimension 1, one can introduce some category C_2 ([13]) of filtered spaces like V that includes *all* components of the adelic complex such as $\mathbb{A}_{12}(D)$ and the space \mathbb{A}_X itself.

We develop the theory of C_2 spaces in section 5.1. In this category, it is possible to define (section 5.2) the spaces of virtual measures $\mu(F(i)|F(j))$ which are crucial for the definition of the spaces of functions and distributions (section 5.3). In section 5.4 we introduce the Fourier transform \mathbf{F} , which preserves the spaces \mathcal{D} and \mathcal{D}' , but interchanges the spaces \mathcal{E} and $\tilde{\mathcal{E}}$. Next section 5.5 contains a study of a central extension of automorphism group of an object V in the category C_2 and its action on the functional spaces such as $\mathcal{D}_{P_0}(V)$ and $\mathcal{D}'_{P_0}(V)$. The definition of central extension goes up to the paper of V. G. Kac and D. H. Peterson, [7], but the definition of an action of group was introduced by M. M. Kapranov for the case of local fields [9]. We then study direct and inverse images (section 5.6), base change rules (section 5.7), the Fourier transform \mathbf{F} (section 5.8), which interchanges direct and inverse images, and, at last, we can prove a generalization of the Poisson formula (section 5.9). It is important that we have two basically different types of this formula (theorems 2 and 3). The formulation includes characteristic functions δ_W of subspaces $W \subset V$ and resembles the formula for dimension 1. It is important that for a class of spaces V (but not for \mathbb{A}_X itself) there exists an invariant measure ($\mathbf{1}_\mu$ or $\mathbf{1}$), defined up to a constant, as an element of $\mathcal{D}'(V)$ or $\mathcal{D}(V)$, an analogue of the classical Haar measure (see section 5.9).

There is also an analytical expression for the intersection number of two divisors based on an adelic approach to the intersection theory, [15, 4]. As a corollary, we get an analytical proof of the (easy part of) Riemann-Roch theorem for divisors on X . This will be exposed in a separate paper.

The main principles of the construction of harmonic analysis were outlined by the second author (see, in particular [18, 17, 19]). The key point here is an observation concerning structure of the classical Bruhat functional spaces on local fields or adelic spaces. Namely, they can be presented as double (projective and/or inductive) limit of functional spaces on finite groups (in particular, on finite-dimensional vector spaces over a finite field). Thus, the construction of harmonic analysis must start with the case of dimension 0 (finite-dimensional vector spaces over a finite field representing a scheme of dimension 0 or finite abelian groups) and then be developed by induction to the higher dimensions. This plan is completely fulfilled in this paper in the cases of dimensions 1 and 2.

An important contribution belongs to M. M. Kapranov [9] who suggested to use a trick from a construction of the Sato Grassmanian in the theory of integrable systems. The trick is to use in the above definition of the spaces $\mathcal{D}_{P_0}(V)$ and $\mathcal{D}'_{P_0}(V)$ the spaces $\mu(P_0/Q)$ of measures instead of $\mu(P/Q)$. Without this trick with the space of measures we cannot define the functional spaces for *all* vector spaces from the category C_2 and, in particular, for the whole adelic space \mathbb{A}_X .

The second author has tried to construct the harmonic analysis in several ways. The most transparent and the easiest one to use is to attach several types of functional spaces to the objects of a *category* of vector spaces arising from some subspaces of the whole adelic space. He worked with a concrete category arising from the subfactors of the adelic space. The key point here is a thorough study of the conditions on homomorphisms of vector spaces that guarantee the existence of direct or inverse images.

Approximately at the same time, the first author independently introduced a notion of C_n structure in the category of vector spaces [13]. With this notion at hand, we can develop the analysis in a very general setting, for any objects of the category C_2 and hopefully for C_n . The crucial point is that the C_n structure exists for the adelic spaces of *any* n -dimensional Noetherian scheme [13, Theorem 2.1]. The principal advantage of this approach is that one can make all the constructions *simultaneously* for local and global cases. The category C_1 contains as a full subcategory the category of linearly locally compact vector spaces (introduced and extensively used by S. Lefschetz [10]) and one can use the classical harmonic analysis under these circumstances.

Note here that the previous publications [18, 17, 19] contained a stupid mistake. Two different types of locally constant functions, $\mathcal{E}(V)$ and $\tilde{\mathcal{E}}(V)$ has not been distinguished (the same was true for distributions).

The present work can certainly be extended, first to the case of spaces arising from arithmetic surfaces. The corresponding category C_2 must include more general abelian groups that appear from local fields of unequal characteristics and the fields of archimedean type. In the last case these are the vector spaces over \mathbb{R} or \mathbb{C} . There are no principal obstacles to do that and we hope to consider this issue in another paper. It seems more difficult but still possible to extend the theory to the

case of higher dimensions, so the category C_n ¹.

In the present work, we develop the harmonic analysis on vector spaces defined either over a local field K or over an adelic ring \mathbb{A} . In classical case of dimension 1, the analysis can be developed on arbitrary varieties (defined either over K , or over \mathbb{A}). This was already done by Bruhat in local case [3]. For arbitrary varieties defined over a two-dimensional local field K , the analysis was constructed by D. Gaitsgory and D. A. Kazhdan in [5] for the purposes of representation theory of reductive groups over the field K (see also arXiv:math/0302174 [math.RT], arXiv:math/0406282 [math.RT], arXiv:math/0409543 [math.RT]). It had been preceded by [8], where harmonic analysis was developed on homogenous spaces such as $G(K)/G(O'_K)$ (introduced in [16]). Note that the construction of harmonic analysis (over \mathbb{A}) is a non-trivial problem even for the case $X = G_m$. This is the subject for further discussion in a separate paper.

Different aspects of the harmonic analysis and its possible applications were exposed by the second author in several talks and lecture series: Münster Universität 1999, 2002, Université Paris VI 2002, 2005, Moscow (Steklov institute) 2002, 2006, Saint-Petersburg (LOMI and Euler Institute) 2003, 2005, Salamanca University 2004, Berlin (Humboldt Universität) 2005, Oberwolfach 2005.

2 Notations

For any vector space V over a field k we will denote by V^* its dual space, i.e.,

$$V^* = \text{Hom}_k(V, k).$$

For a vector k -subspace $H \subset V$ we denote by H^\perp the following k -subspace in the space V^* :

$$H^\perp \stackrel{\text{def}}{=} \{u \in V^* : u(H) = 0\}.$$

By $\mathcal{F}(V)$ we denote the space of all functions on V with values in the field \mathbf{C} .

For any $a \in V$, for any $f \in \mathcal{F}(V)$ by $T_a(f)$ we denote the following function from $\mathcal{F}(V)$:

$$T_a(f)(v) \stackrel{\text{def}}{=} f(v + a), \quad v \in V.$$

For any $f \in \mathcal{F}(V)$ by \check{f} we denote the following function from $\mathcal{F}(V)$:

$$\check{f}(v) \stackrel{\text{def}}{=} f(-v), \quad v \in V.$$

For a finite set A we denote by $\sharp A$ the number of elements in the set A . By $\text{supp}(f)$ we denote the support of a function f .

We fix a finite field \mathbb{F}_q and a nontrivial character $\psi : \mathbb{F}_q \rightarrow \mathbf{C}^*$.

¹In the beginning of July 2007, our paper was put on the net arXiv:0707.1766 [math.AG]. Then, there appeared a paper by A. Deitmar arXiv:0708.0322 [math.NT], where he claimed that he constructed harmonic analysis for all dimensions n . Unfortunately, the first versions of his work contained serious mistakes that were pointed out to the author in our letters. The latest version of his work contains only a construction of the spaces like \mathcal{E} , which is in fact a rather trivial generalization of our work.

3 Zero-dimensional case

Let V be a finite-dimensional vector space over the field \mathbb{F}_q .

Then any element $f \in \mathcal{F}(V)$ has the following unique presentation:

$$f = \sum_{v \in V} a_v \cdot \delta_v,$$

where $a_v \in \mathbf{C}$, and the functions $\delta_v \in \mathcal{F}(V)$ are defined by the following rule

$$\delta_v(w) = \begin{cases} 1, & \text{if } w = v \\ 0, & \text{if } w \neq v. \end{cases}$$

On the space $\mathcal{F}(V)$ we have the following pairing $\langle, \rangle_V: \mathcal{F}(V) \times \mathcal{F}(V) \longrightarrow \mathbf{C}$:

$$\langle f, g \rangle_V = \sum_{v \in V} f(v) \cdot g(v).$$

We have the following obvious proposition.

Proposition 1 *The pairing \langle, \rangle_V is a nondegenerate symmetric pairing on the finite-dimensional \mathbf{C} -vector space $\mathcal{F}(V)$, and therefore this pairing determines a canonical isomorphism between the vector space $\mathcal{F}(V)$ and the dual vector space $\mathcal{F}(V)^*$.*

Let $\pi: V \rightarrow W$ be a linear map between finite-dimensional vector spaces over the field \mathbb{F}_q . We define the direct image $\pi_*: \mathcal{F}(V) \rightarrow \mathcal{F}(W)$ as the following map:

$$\pi_*(f)(w) = \begin{cases} 0 & , \text{ if } w \notin \pi(V) \\ \sum_{\substack{v \in V \\ \pi(v)=w}} f(v) & , \text{ if } w \in \pi(V), \end{cases}$$

where $f \in \mathcal{F}(V)$, $w \in W$.

We define the inverse image $\pi^*: \mathcal{F}(W) \rightarrow \mathcal{F}(V)$ as the following map:

$$\pi^*(g)(v) = f(\pi(v)) \quad , \text{ where } g \in \mathcal{F}(W), \quad v \in V.$$

There is the following proposition, which follows easily from the definitions.

Proposition 2 *The maps π_* and π^* are conjugate maps with respect to the pairings \langle, \rangle_V and \langle, \rangle_W on the spaces $\mathcal{F}(V)$ and $\mathcal{F}(W)$, i.e.,*

$$\langle \pi^*(g), f \rangle_V = \langle g, \pi_*(f) \rangle_W$$

for any $f \in \mathcal{F}(V)$, $g \in \mathcal{F}(W)$.

There is also the following proposition, which also easily follows from the definitions.

Proposition 3 *The following statements are satisfied.*

1. Let $\pi_1 : V_1 \rightarrow V_2$ and $\pi_2 : V_2 \rightarrow V_3$ are linear maps between the vector spaces over the field \mathbb{F}_q , then

$$(\pi_2)_*(\pi_1)_*(f) = (\pi_2\pi_1)_*(f) \quad \text{for any } f \in \mathcal{F}(V_1),$$

$$\text{and } \pi_1^*\pi_2^*(g) = (\pi_2\pi_1)^*(g) \quad \text{for any } g \in \mathcal{F}(V_3).$$

2. Let V, W, S are finite-dimensional vector spaces over the field \mathbb{F}_q . Let $\pi : V \rightarrow S, \alpha : W \rightarrow S$ are linear maps between these spaces. We consider the following cartesian diagram of vector spaces:

$$\begin{array}{ccc} V \times_S W & \xrightarrow{\pi_W} & W \\ \alpha_V \downarrow & & \downarrow \alpha \\ V & \xrightarrow{\pi} & S \end{array}$$

Then we have the following base change rule

$$\pi^*\alpha_*(h) = (\alpha_V)_*\pi_W^*(h) \quad \text{for any } h \in \mathcal{F}(W).$$

For a finite-dimensional vector space V over the field \mathbb{F}_q we define the Fourier transform $\mathbf{F} : \mathcal{F}(V) \rightarrow \mathcal{F}(V^*)$ by the following rule:

$$\mathbf{F}(f)(u) = \sum_{v \in V} f(v) \cdot \overline{\psi(u(v))},$$

where $u \in V^*, f \in \mathcal{F}(V)$. We consider the following example.

Example 1 Let $H \subset V$ be a vector subspace. We consider the following function $\delta_H = \sum_{v \in H} \delta_v$ from $\mathcal{F}(V)$. Then it is easy to see by direct calculation that

$$\mathbf{F}(\delta_H) = (\sharp H) \cdot \delta_{H^\perp}.$$

Remark 1 Example 1 is the *Poisson formula* for 0-dimensional case.

We have the following proposition, which collects the properties of the Fourier transform \mathbf{F} .

Proposition 4 *The following statements are satisfied.*

1. \mathbf{F} is an isomorphism of \mathbf{C} -vector spaces.
2. $\mathbf{F} \circ \mathbf{F}(f) = \sharp V \cdot \check{f}$ for any $f \in \mathcal{F}(V)$.
3. $\langle \mathbf{F}(f), g \rangle_{V^*} = \langle f, \mathbf{F}(g) \rangle_V$ for any $f \in \mathcal{F}(V), g \in \mathcal{F}(V^*)$.

4. Let $\pi : V \rightarrow W$ be a linear map between the finite-dimensional vector spaces over the field \mathbb{F}_q . Let $\pi' : W^* \rightarrow V^*$ be the dual map. Then the following diagrams are commutative:

$$\begin{array}{ccc} \mathcal{F}(V) & \xrightarrow{\pi_*} & \mathcal{F}(W) \\ \mathbf{F} \downarrow & & \downarrow \mathbf{F} \\ \mathcal{F}(V^*) & \xrightarrow{(\pi')^*} & \mathcal{F}(W^*) \end{array} \quad (1)$$

$$\begin{array}{ccc} \mathcal{F}(W) & \xrightarrow{\pi^*} & \mathcal{F}(V) \\ \frac{1}{\sharp W} \cdot \mathbf{F} \downarrow & & \downarrow \mathbf{F} \cdot \frac{1}{\sharp V} \\ \mathcal{F}(W^*) & \xrightarrow{(\pi')^*} & \mathcal{F}(V^*) \end{array} \quad (2)$$

Proof . Statements 2, 3 and diagram (1) follow by direct calculations on functions δ_v . Now statement 1 follows from statement 2. Diagram (2) follows from diagram (1) and statement 2.

4 One-dimensional case

4.1 C_0 and C_1 -spaces.

Here we recall the definitions from [13] about C_0 and C_1 -spaces.

By definition, the category C_0 is the category of finite-dimensional vector spaces over a field k with morphisms coming from k -linear maps between vector spaces.

By definition, an admissible triple of finite-dimensional vector spaces is an exact triple of this vector spaces.

We say that (I, F, V) is a filtered k -vector space, if

- (i) V is a vector space over the field k ,
- (ii) I is a partially ordered set, such that for any $i, j \in I$ there are $k, l \in I$ with $k \leq i \leq l$ and $k \leq j \leq l$,
- (iii) F is a function from I to the set of k -vector subspaces of V such that if $i \leq j$ are any from I , then $F(i) \subset F(j)$,
- (iv) $\bigcap_{i \in I} F(i) = 0$ and $\bigcup_{i \in I} F(i) = V$.

We say that a filtered vector space (I_1, F_1, V) *dominates* another filtered vector space (I_2, F_2, V) when there is an order-preserving function $\phi : I_2 \rightarrow I_1$ such that

- (i) for any $i \in I_2$ we have $F_1(\phi(i)) = F_2(i)$,
- (ii) for any $j \in I_1$ there are $i_1, i_2 \in I_2$ such that $\phi(i_1) \leq j \leq \phi(i_2)$.

By definition, objects of the category C_1 , i.e. $\text{Ob}(C_1)$, are filtered k -vector spaces (I, F, V) such that for any $i \leq j \in I$ the k -vector space $F(j)/F(i)$ is finite-dimensional over the field k .

Let $E_1 = (I_1, F_1, V_1)$ and $E_2 = (I_2, F_2, V_2)$ be from $\text{Ob}(C_1)$. Then, by definition, a set $\text{Mor}_{C_1}(E_1, E_2)$ consists of elements $A \in \text{Hom}_k(V_1, V_2)$ under the following condition: for any $j \in I_2$ there is an $i \in I_1$ such that $A(F_1(i)) \subset F_2(j)$.

Let $E_1 = (I_1, F_1, V_1)$, $E_2 = (I_2, F_2, V_2)$ and $E_3 = (I_3, F_3, V_3)$ be from $\text{Ob}(C_1)$. Then we say that

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0 \quad (3)$$

is an *admissible triple* from C_1 when the following conditions are satisfied:

(i)

$$0 \longrightarrow V_1 \xrightarrow{\alpha} V_2 \xrightarrow{\beta} V_3 \longrightarrow 0$$

is an exact triple of k -vector spaces,

(ii) the filtration (I_1, F_1, V_1) dominates the filtration (I_2, F'_1, V_1) , where $F'_1(i) = F_2(i) \cap V_1$ for any $i \in I_2$,

(iii) the filtration (I_3, F_3, V_3) dominates the filtration (I_2, F'_3, V_3) , where $F'_3(i) = F_2(i)/F_2(i) \cap V_1$ for any $i \in I_2$,

(iv) for any $i \leq j \in I_2$

$$0 \longrightarrow \frac{F'_1(j)}{F'_1(i)} \longrightarrow \frac{F_2(j)}{F_2(i)} \longrightarrow \frac{F'_3(j)}{F'_3(i)} \longrightarrow 0$$

is an exact triple of finite-dimensional k -vector spaces.

We say that α from an admissible triple (3) is an *admissible monomorphism*, and β from an admissible triple (3) is an *admissible epimorphism*.

Remark 2 Let $E_2 = (I_2, F_2, V_2)$ be a C_1 -space over the field k . Let

$$0 \longrightarrow V_1 \xrightarrow{\alpha} V_2 \xrightarrow{\beta} V_3 \longrightarrow 0 \quad (4)$$

be an exact triple of k -vector spaces. We define on the spaces V_1 and V_3 the structures of filtered spaces (I_i, F_i, V_i) , $i = 1, 3$, where $I_1 = I_3 = I_2$, and $F_1(i) \stackrel{\text{def}}{=} F_2(i) \cap V_1$, $F_3(i) \stackrel{\text{def}}{=} \beta(F_2(i))$ for $i \in I_2$.

Then the filtered space $E_1 = (I_1, F_1, V_1)$ will be always a C_1 -space. But filtered space $E_3 = (I_3, F_3, V_3)$ will be a C_1 -space, if and only if the following condition holds:

(*) for any $x \in V_2 \setminus V_1$ there exists $i \in I_2$ such that $(x + F_2(i)) \cap V_1 = \emptyset$.

Moreover, under condition (*), exact triple (4) gives the admissible triple

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0,$$

i.e. α will be an admissible monomorphism, β will be an admissible epimorphism.

For example, we consider $k((t))$ as a C_1 -space with filtration, given by all powers of maximal ideal of the ring $k[[t]]$. Then k -subspace $k[t]$ with induced filtration will be a C_1 -space, but the embedding $k[t] \subset k((t))$ will not be an admissible monomorphism. And embedding $k[t^{-1}] \subset k((t))$ will be an admissible monomorphism (with the induced filtration).

Let $E = (I, F, V)$ be a C_1 -space. We define a C_1 -space $\check{E} = (I^0, F^0, \check{V})$ by the following way. The space $\check{V} \subset V^*$ is defined as

$$\check{V} \stackrel{\text{def}}{=} \bigcup_{i \in I} F(i)^\perp. \quad (5)$$

The set I^0 is a partially ordered set, which has the same set as I , but with the inverse order then I . For $j \in I^0$ the subspace $F^0(j) \stackrel{\text{def}}{=} F(j)^\perp \subset \check{V}$.

We note also that

$$F(j)^\perp = (V/F(j))^* = \lim_{\substack{\leftarrow \\ i \geq j}} (F(i)/F(j))^*.$$

If $E_1, E_2 \in \text{Ob}(C_1)$ and $\theta \in \text{Mor}_{C_1}(E_1, E_2)$. Then there is canonically $\check{\theta} \in \text{Mor}_{C_1}(\check{E}_2, \check{E}_1)$. If

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

is an admissible triple of C_1 -spaces, where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. Then there is canonically the following admissible triple of C_1 -spaces:

$$0 \longrightarrow \check{E}_3 \xrightarrow{\check{\beta}} \check{E}_2 \xrightarrow{\check{\alpha}} \check{E}_1 \longrightarrow 0.$$

Lemma 1 *Let*

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_1 -spaces.

1. *Let $D \in \text{Ob}(C_1)$, and $\gamma \in \text{Mor}_{C_1}(D, E_3)$. Then there is the following admissible triple of C_1 -spaces*

$$0 \longrightarrow E_1 \xrightarrow{\gamma\alpha} E_2 \times_{E_3} D \xrightarrow{\gamma\beta} D \longrightarrow 0 \quad (6)$$

and $\beta_\gamma \in \text{Mor}_{C_1}(E_2 \times_{E_3} D, E_2)$ such that the following diagram is commutative:

$$\begin{array}{ccccccc} 0 & \longrightarrow & E_1 & \xrightarrow{\gamma\alpha} & E_2 \times_{E_3} D & \xrightarrow{\gamma\beta} & D \longrightarrow 0 \\ & & \parallel & & \downarrow \beta_\gamma & & \downarrow \gamma \\ 0 & \longrightarrow & E_1 & \xrightarrow{\alpha} & E_2 & \xrightarrow{\beta} & E_3 \longrightarrow 0 \end{array} \quad (7)$$

2. *Let $A \in \text{Ob}(C_1)$, and $\theta \in \text{Mor}_{C_1}(E_1, A)$. Then there is the following admissible triple of C_1 -spaces*

$$0 \longrightarrow A \xrightarrow{\theta\alpha} A \amalg_{E_1} E_2 \xrightarrow{\theta\beta} E_3 \longrightarrow 0 \quad (8)$$

and $\alpha_\theta \in \text{Mor}_{C_1}(E_2, A \amalg_{E_1} E_2)$ such that the following diagram is commutative:

$$\begin{array}{ccccccccc}
0 & \longrightarrow & E_1 & \xrightarrow{\alpha} & E_2 & \xrightarrow{\beta} & E_3 & \longrightarrow & 0 \\
& & \downarrow \theta & & \downarrow \alpha_\theta & & \parallel & & \\
0 & \longrightarrow & A & \xrightarrow{\theta_\alpha} & A \amalg_{E_1} E_2 & \xrightarrow{\theta_\beta} & E_3 & \longrightarrow & 0
\end{array} \tag{9}$$

Proof . We prove the first statement of the lemma. Let $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$, and $D = (K, H, T)$.

We construct $E_2 \times_{E_3} D = (J, G, W) \in \text{Ob}(C_1)$ in the following way. We define a k -vector space

$$W = V_2 \times_{V_3} T \stackrel{\text{def}}{=} \{(e, d) \in V_2 \times T \text{ such that } \beta(e) = \gamma(d)\}.$$

We define a partially ordered set

$$J = \{(i, j) \in I_2 \times K \text{ such that } \gamma(H(j)) \subset \beta(F_2(i))\},$$

where $(i_1, j_1) \leq (i_2, j_2)$ iff $i_1 \leq i_2$ and $j_1 \leq j_2$. We define a function G from J to the set of k -vector subspaces of W as

$$G((i, j)) = (F_2(i) \times H(j)) \cap W,$$

where the intersection is inside $V_2 \times T$. Then $E_2 \times_{E_3} D$ is well defined as a C_1 -space, and the maps β_γ and γ_β are projections.

Let $(i_1, j_1), (i_2, j_2) \in J$, and $(i_1, j_1) \leq (i_2, j_2)$. Then we have the following commutative diagram of finite-dimensional k -vector spaces:

$$\begin{array}{ccccccc}
0 & \longrightarrow & \frac{F_2(i_2) \cap V_1}{F_2(i_1) \cap V_1} & \longrightarrow & \frac{G((i_2, j_2))}{G((i_1, j_1))} & \longrightarrow & \frac{H(j_2)}{H(j_1)} \longrightarrow 0 \\
& & \parallel & & \downarrow & & \downarrow \\
0 & \longrightarrow & \frac{F_2(i_2) \cap V_1}{F_2(i_1) \cap V_1} & \longrightarrow & \frac{F_2(i_2)}{F_2(i_1)} & \longrightarrow & \frac{F_2(i_2)/(F_2(i_2) \cap V_1)}{F_2(i_1)/(F_2(i_1) \cap V_1)} \longrightarrow 0
\end{array}$$

where

$$\frac{G((i_2, j_2))}{G((i_1, j_1))} = \frac{F_2(i_2)}{F_2(i_1)} \times_{\frac{F_2(i_2)/(F_2(i_2) \cap V_1)}{F_2(i_1)/(F_2(i_1) \cap V_1)}} \frac{H(j_2)}{H(j_1)}.$$

From this commutative diagram we obtain that triple (6) is an admissible triple of C_1 -spaces, and diagram (7) is a commutative diagram. The first statement of the lemma is proved.

We prove the second statement of the lemma, which is the dual statement to the first statement of the lemma. Let $A = (K', H', T')$.

We construct $A \amalg_{E_1} E_2 = (J', G', W') \in \text{Ob}(C_1)$ in the following way. We remark that $T' \amalg V_2 = T' \times V_2$. We define a k -vector space

$$W' = T' \amalg_{V_1} V_2 \stackrel{\text{def}}{=} (T' \times V_2)/E \quad , \quad \text{where} \quad E = (\theta \times \alpha)V_1.$$

We define a partially ordered set

$$J' = \{(i, j) \in K' \times I_2 : \theta(F_2(j) \cap V_1) \subset H'(i)\},$$

where $(i_1, j_1) \leq (i_2, j_2)$ iff $i_1 \leq i_2$ and $j_1 \leq j_2$. We define a function G' from J' to the set of k -vector subspaces of W' as

$$G'((i, j)) = ((H'(i) \times F_2(j)) + E/E) \subset W'.$$

Then $A \amalg_{E_1} E_2$ is well defined as a C_1 -space.

Let $(i_1, j_1), (i_2, j_2) \in J'$, and $(i_1, j_1) \leq (i_2, j_2)$. Then we have the following commutative diagram of finite-dimensional k -vector spaces:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \frac{F_2(j_2) \cap V_1}{F_2(j_1) \cap V_1} & \longrightarrow & \frac{F_2(j_2)}{F_2(j_1)} & \longrightarrow & \frac{F_2(j_2)/(F_2(j_2) \cap V_1)}{F_2(j_1)/(F_2(j_1) \cap V_1)} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \frac{H'(i_2)}{H'(i_1)} & \longrightarrow & \frac{G'((i_2, j_2))}{G'((i_1, j_1))} & \longrightarrow & \frac{F_2(j_2)/(F_2(j_2) \cap V_1)}{F_2(j_1)/(F_2(j_1) \cap V_1)} \longrightarrow 0 \end{array}$$

where

$$\frac{G'((i_2, j_2))}{G'((i_1, j_1))} = \frac{H'(i_2)}{H'(i_1)} \amalg_{\frac{F_2(j_2) \cap V_1}{F_2(j_1) \cap V_1}} \frac{F_2(j_2)}{F_2(j_1)}.$$

From this commutative diagram we obtain that triple (8) is an admissible triple of C_1 -spaces, and diagram (9) is a commutative diagram. The second statement of the lemma is proved.

Definition 1 Let $(I, F, V), (J, G, V) \in \text{Ob}(C_1)$. We say that there is an equivalence $(I, F, V) \sim (J, G, V)$ iff there is a collection of C_1 -spaces (I_l, F_l, V) , $1 \leq l \leq n$ such that

- (i) $I_1 = I$, $F_1 = F$, $I_n = J$, $F_n = G$,
- (ii) for any $1 \leq l \leq n-1$ either (I_l, F_l, V) dominates (I_{l+1}, F_{l+1}, V) , or (I_{l+1}, F_{l+1}, V) dominates (I_l, F_l, V) .

Let $\theta : E \longrightarrow D$ be an admissible epimorphism of C_1 -spaces. Let $\beta : B \longrightarrow D$ be an admissible monomorphism of C_1 -spaces. Then from construction (see the proof of lemma 1) we have that $\beta_\theta : E \times_D B \longrightarrow B$ is an admissible epimorphism of C_1 -spaces, and $\theta_\beta : E \times_D B \longrightarrow E$ is an admissible monomorphism.

Moreover, Let $N = E \times_D B$. From construction we have that $D \sim B \amalg_N E$. We have the admissible epimorphism $\check{\theta}_\beta : \check{E} \longrightarrow \check{N}$, and the admissible monomorphism $\check{\beta}_\theta : \check{B} \longrightarrow \check{N}$. From construction (see the proof of lemma 1) we have that $\check{D} \sim \check{E} \times_{\check{N}} \check{B}$.

Definition 2 We say that a C_1 -space (I_1, F_1, V_1) is a compact C_1 -space iff there is an $i \in I_1$ such that $F_1(i) = V_1$. We say that a C_1 -space (I_2, F_2, V_2) is a discrete C_1 -space iff there is an $j \in I_2$ such that $F_2(j) = \{0\}$.

It follows from definition that if E_1 is a compact C_1 -space, then \check{E}_1 is a discrete C_1 -space. If E_2 is a discrete C_1 -space, then \check{E}_2 is a compact C_1 -space.

Definition 3 We say that a C_1 -space (I, F, V) is a complete C_1 -space iff

$$V = \lim_{\substack{\rightarrow \\ i \in I}} \lim_{\substack{\leftarrow \\ j \leq i}} F(i)/F(j).$$

It follows from definitions that E is a complete C_1 -space iff $\check{E} = E$.

4.2 Spaces of functions and distributions.

We recall the basic definitions from [18] and [19].

Let $E = (I, F, V)$ be a C_1 -space over the field \mathbb{F}_q . For any $i, j, k \in I$ such that $i \leq j \leq k$ we have the embedding map

$$\alpha_{ijk} : F(j)/F(i) \hookrightarrow F(k)/F(i)$$

and the surjection map

$$\pi_{ijk} : F(k)/F(i) \twoheadrightarrow F(k)/F(j)$$

of finite-dimensional vector spaces over the field \mathbb{F}_q .

Then according to section 3 we have 4 maps:

$$\begin{aligned} \alpha_{ijk}^* & : \mathcal{F}(F(k)/F(i)) \longrightarrow \mathcal{F}(F(j)/F(i)), \\ (\alpha_{ijk})_* & : \mathcal{F}(F(j)/F(i)) \longrightarrow \mathcal{F}(F(k)/F(i)), \\ \pi_{ijk}^* & : \mathcal{F}(F(k)/F(j)) \longrightarrow \mathcal{F}(F(k)/F(i)), \\ (\pi_{ijk})_* & : \mathcal{F}(F(k)/F(i)) \longrightarrow \mathcal{F}(F(k)/F(j)). \end{aligned}$$

According to the maps described, we consider now all possible combinations of projective and inductive limits of spaces $\mathcal{F}(F(j)/F(i))$ with respect to the indices $i, j \in I$, $i \leq j$. We define

$$\mathcal{D}(E) \stackrel{\text{def}}{=} \lim_{\substack{\rightarrow \\ j}} \lim_{\substack{\rightarrow \\ i}} \mathcal{F}(F(j)/F(i)) = \lim_{\substack{\rightarrow \\ i}} \lim_{\substack{\rightarrow \\ j}} \mathcal{F}(F(j)/F(i)) \quad (10)$$

with respect to the maps π_{ijk}^* and $(\alpha_{ijk})_*$;

$$\mathcal{D}'(E) \stackrel{\text{def}}{=} \lim_{\substack{\leftarrow \\ j}} \lim_{\substack{\leftarrow \\ i}} \mathcal{F}(F(j)/F(i)) = \lim_{\substack{\leftarrow \\ i}} \lim_{\substack{\leftarrow \\ j}} \mathcal{F}(F(j)/F(i)) \quad (11)$$

with respect to the maps $(\pi_{ijk})_*$ and α_{ijk}^* ;

$$\mathcal{E}(E) \stackrel{\text{def}}{=} \lim_{\substack{\leftarrow \\ j}} \lim_{\substack{\rightarrow \\ i}} \mathcal{F}(F(j)/F(i)) \quad (12)$$

with respect to the maps α_{ijk}^* and π_{ijk}^* ;

$$\tilde{\mathcal{E}}(E) \stackrel{\text{def}}{=} \lim_{\substack{\rightarrow \\ i}} \lim_{\substack{\leftarrow \\ j}} \mathcal{F}(F(j)/F(i)) \quad (13)$$

with respect to the maps π_{ijk}^* and α_{ijk}^* ;

$$\mathcal{E}'(E) \stackrel{\text{def}}{=} \lim_{\substack{\rightarrow \\ j}} \lim_{\substack{\leftarrow \\ i}} \mathcal{F}(F(j)/F(i)) \quad (14)$$

with respect to the maps $(\alpha_{ijk})_*$ and $(\pi_{ijk})_*$;

$$\tilde{\mathcal{E}}'(E) \stackrel{\text{def}}{=} \lim_{\substack{\leftarrow \\ i}} \lim_{\substack{\rightarrow \\ j}} \mathcal{F}(F(j)/F(i)) \quad (15)$$

with respect to the maps $(\pi_{ijk})_*$ and $(\alpha_{ijk})_*$.

Due to proposition 3 we have that \mathbf{C} -vector spaces $\mathcal{D}(E)$, $\mathcal{D}'(E)$, $\mathcal{E}(E)$, $\mathcal{E}(\tilde{E})$, $\mathcal{E}'(E)$, $\tilde{\mathcal{E}}'(E)$ are well-defined.

Since $\alpha_{ijk}^* \circ (\alpha_{ijk})_*$ is the identity map for any $i \leq j \leq k \in I$, we have the following canonical embeddings

$$\mathcal{D}(E) \hookrightarrow \tilde{\mathcal{E}}(E) \quad (16)$$

$$\tilde{\mathcal{E}}'(E) \hookrightarrow \mathcal{D}'(E). \quad (17)$$

We have also the following canonical embeddings

$$\tilde{\mathcal{E}}(E) \hookrightarrow \mathcal{E}(E) \quad (18)$$

$$\mathcal{E}'(E) \hookrightarrow \tilde{\mathcal{E}}'(E). \quad (19)$$

Now we clarify the functional sense of spaces $\mathcal{D}(E)$, $\mathcal{D}'(E)$, $\mathcal{E}(E)$, $\tilde{\mathcal{E}}(E)$, $\mathcal{E}'(E)$.

Proposition 5 *The following statements are satisfied.*

1. *The space $\mathcal{D}(E) \subset \mathcal{F}(V)$ such that:*

$$f \in \mathcal{D}(E) \quad \text{iff there are } i \leq j \in I \quad \text{such that}$$

$$f(v) = 0 \quad \text{for any } v \notin F(j) \quad \text{and} \quad T_w(f) = f \quad \text{for any } w \in F(i),$$

i.e., the space $\mathcal{D}(E)$ consists of all locally constant functions with compact support on the space V .

2. *We suppose that $E = (I, F, V)$ is a complete C_1 -space. Then the space $\mathcal{E}(E) \subset \mathcal{F}(V)$ such that:*

$$f \in \mathcal{E}(E) \quad \text{iff for any } v \in V \quad \text{there is } i \in I \quad (\text{which depends on } v)$$

$$\text{such that } f(v+w) = f(v) \quad \text{for any } w \in F(i),$$

i.e., the space $\mathcal{E}(E)$ consists of all locally constant functions on the space V .

3. The space $\tilde{\mathcal{E}}(E) \subset \mathcal{F}(V)$ such that:

$$f \in \tilde{\mathcal{E}}(E) \text{ iff there is } i \in I \text{ such that } T_w(f) = f \text{ for any } w \in F(i).$$

4. The space $\mathcal{D}'(E) = \mathcal{D}(E)^*$, i.e., the space $\mathcal{D}'(E)$ consists of distributions on the space V .

Proof . The first, the second, and the third statement of this proposition are just reformulation of formulas (10), (12), (13). In the second statement we demand that V is a complete C_1 -space, because then for any locally constant function $f \in \mathcal{F}(V)$ for any $j \in I$ there is an $i \in I$ such that $f(v + w) = f(v)$ for any $v \in F(j)$, $w \in F(i)$.

To prove the fourth statement we remark that

$$\mathcal{D}(E)^* = \lim_{\leftarrow j} \lim_{\leftarrow i} \mathcal{F}(F(j)/F(i))^*,$$

because the dual space to the space given by inductive limits of vector spaces is given by projective limits of dual vector spaces. Now we apply propositions 1 and 2 to obtain that $\mathcal{D}'(E) = \mathcal{D}(E)^*$.

We can not speak about the value of distribution $G \in \mathcal{D}'(E)$ in the point $v \in V$. But we can say, whether G is equal to zero or not in the point $v \in V$. We have the following definition.

Definition 4 Let $G \in \mathcal{D}'(E)$.

1. For $v \in V$ we say $G(v) = 0$ iff there is $i \in I$ such that $G(f) = 0$ for any $f \in \mathcal{D}(V)$ with $\text{supp } f \subset \{v + F(i)\}$.

2.

$$\text{supp } G \stackrel{\text{def}}{=} \{v \in V : G(v) \neq 0\}.$$

Due to definition 4 we have the following proposition.

Proposition 6 Let $E = (I, F, V)$ be a complete C_1 -space. For $G \in \mathcal{D}'(E)$ we have that

$$G \in \mathcal{E}'(E) \text{ iff there is } j \in I \text{ such that } \text{supp } G \subset F(j),$$

i.e., the space $\mathcal{E}'(E)$ consists of all distributions with compact support on the space V .

Proof follows from the forth statement of proposition 5 and formula (14).

Remark 3 Using formulas (10) – (15) we obtain that

$$\mathcal{E}(E) = \tilde{\mathcal{E}}(E) = \mathcal{D}(E) \quad \text{and} \quad \mathcal{E}'(E) = \tilde{\mathcal{E}}'(E) = \mathcal{D}'(E)$$

for any compact C_1 -space $E = (I, F, V)$.

Using formulas (10) – (15) we obtain that

$$\tilde{\mathcal{E}}'(E) = \mathcal{E}'(E) = \mathcal{D}(E) \quad \text{and} \quad \tilde{\mathcal{E}}(E) = \mathcal{E}(E) = \mathcal{D}'(E)$$

for any discrete C_1 -space $E = (I, F, V)$.

Remark 4 Let $\{B_k\}$ be a projective system of vector spaces with surjective transition maps. Then we have a canonical nondegenerate pairing between vector spaces

$$\varprojlim_k B_k \quad \text{and} \quad \varinjlim_k B_k^*.$$

Using it and formulas (12), (14), (15), (13) we have canonical nondegenerate pairings

$$\begin{aligned} \langle, \rangle_{\mathcal{E}(E)} &: \mathcal{E}(E) \times \mathcal{E}'(E) \longrightarrow \mathbf{C}, \\ \langle, \rangle_{\tilde{\mathcal{E}}(E)} &: \tilde{\mathcal{E}}'(E) \times \tilde{\mathcal{E}}(E) \longrightarrow \mathbf{C}. \end{aligned}$$

These pairings, being restricted to $\mathcal{D}(E) \times \mathcal{E}'(E)$ and to $\tilde{\mathcal{E}}'(E) \times \mathcal{D}(E)$ correspondingly, coincide with the pairings restricted from the pairing between $\mathcal{D}(E)$ and $\mathcal{D}'(E)$ (see statement 4 of proposition 5).

4.3 Invariant measures and integral

Let $E = (I, F, V)$ be a C_1 -space over the field \mathbb{F}_q . For any $G \in \mathcal{D}'(E)$, for any $a \in V$ we denote by $T_a(G)$ the following distribution from $\mathcal{D}'(E)$:

$$T_a(G)(f) \stackrel{\text{def}}{=} G(T_{-a}(f)), \quad f \in \mathcal{D}(E).$$

We define

$$\mathcal{D}'(E)^V \stackrel{\text{def}}{=} \{G \in \mathcal{D}'(E) \text{ such that } T_a(G) = G \text{ for any } a \in V\}.$$

We have the following proposition.

Proposition 7 $\mathcal{D}'(E)^V$ is a 1-dimensional vector space over the field \mathbf{C} .

Proof. We note that the space $\mathcal{D}(E)$ is generated by functions $T_a(\delta_{F(i)})$, $a \in V$, $i \in I$, where the function

$$\delta_{F(i)}(v) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } v \in F(i) \\ 0, & \text{if } v \notin F(i). \end{cases}$$

Besides, we have for any $i \leq j \in I$

$$\delta_{F(j)} = \sum_{b \in F(j)/F(i)} T_{a_b}(\delta_{F(i)}),$$

where $a_b \in F(j)$ is some lift of $b \in F(j)/F(i)$.

Therefore for fixed element $i \in I$ it is enough to define $G \in \mathcal{D}'(E)^V$ only on $\delta_{F(i)} \in \mathcal{D}(E)$. The space of these values is a 1-dimensional vector space over the field \mathbf{C} .

We denote the 1-dimensional \mathbf{C} -vector space $\mathcal{D}'(E)^V$ by $\mu(E)$ and call it the space of *invariant measures* on E . For any $\mu \in \mu(E)$ and for any $f \in \mathcal{D}(E)$ we denote

$$\int_V f d\mu \stackrel{\text{def}}{=} \mu(f).$$

For any $i \in I$ we denote

$$\mu(F(i)) \stackrel{\text{def}}{=} \mu(\delta_{F(i)}) = \int_V \delta_{F(i)} d\mu.$$

We note that the \mathbf{C} -vector space $\mathcal{F}(V)$ is a \mathbf{C} -algebra, i.e., for any $f, g \in \mathcal{F}(V)$ we have $f \cdot g(v) \stackrel{\text{def}}{=} f(v) \cdot g(v)$, where $v \in V$. From definitions of subspaces $\mathcal{E}(E) \subset \mathcal{F}(V)$ and $\mathcal{D}(E) \subset \mathcal{F}(V)$ we have that

$$\mathcal{E}(E) \cdot \mathcal{D}(E) \subset \mathcal{D}(E). \quad (20)$$

Therefore for fixed $\mu \in \mu(E)$ it is well-defined the following map

$$\begin{aligned} \mathbf{I}_\mu &: \mathcal{E}(E) \longrightarrow \mathcal{D}'(E) \\ \mathbf{I}_\mu(f)(g) &\stackrel{\text{def}}{=} \int_V f \cdot g d\mu, \quad f \in \mathcal{E}(E), \quad g \in \mathcal{D}(E). \end{aligned}$$

We have the following proposition.

Proposition 8 *Let $\mu \in \mu(E)$, $\mu \neq 0$. There is the following 6-edges commutative diagram of embeddings:*

$$\begin{array}{ccccc} & & \mathcal{D}'(E) & & \\ & \nearrow \mathbf{I}_\mu & & \nwarrow & \\ \mathcal{E}(E) & & & & \tilde{\mathcal{E}}'(E) \\ \uparrow & & & & \uparrow \\ \tilde{\mathcal{E}}(E) & & & & \mathcal{E}'(E) \\ & \nwarrow & & \nearrow \mathbf{I}_\mu & \\ & & \mathcal{D}(E) & & \end{array} \quad (21)$$

Proof . Most edges are embeddings which given by formulas (16)-(19).

We proof that the map $\mathbf{I}_\mu : \mathcal{E}(E) \rightarrow \mathcal{D}'(E)$ is an embedding. Indeed, let $f \in \mathcal{E}(E)$, $f \neq 0$. Then there is $v \in V$ such that $f(v) \neq 0$. By definition of the space $\mathcal{E}(E)$ there is $i \in I$ such that $f(v+w) = f(v)$ for any $w \in F(i)$. Then $\mathbf{I}_\mu(f)(T_v(\delta_{F(i)})) = f(v) \cdot \mu(F(i)) \neq 0$. Therefore $\mathbf{I}_\mu(f) \neq 0$.

Now we prove that $\mathbf{I}_\mu(\mathcal{D}(E)) \subset \mathcal{E}'(E)$. Indeed, let $f \in \mathcal{D}(E)$, $\text{supp } f \subset F(j)$ for some $j \in I$. Then $\mathbf{I}_\mu(f) \in \varprojlim_{i \leq j} \mathcal{F}(F(j)/F(i))$, because $\mathbf{I}_\mu(f)(g) = \mathbf{I}_\mu(f)(g|_{F(j)})$ for

any $g \in \mathcal{D}(E)$. (Here $g|_{F(j)}$ is the function restricted to $F(j)$ and then extended to V by 0 outside $F(j)$). Therefore $\mathbf{I}_\mu(f) \in \mathcal{E}'(E)$.

Remark 5 Let $f \in \mathcal{E}(E)$, $G \in \mathcal{D}'(E)$. Then we define $f \cdot G \in \mathcal{D}'(E)$ by the following rule:

$$f \cdot G(g) \stackrel{\text{def}}{=} G(f \cdot g), \quad g \in \mathcal{D}(E).$$

From definitions we have for any $f_1, f_2 \in \mathcal{E}(E)$

$$\mathbf{I}_\mu(f_1 \cdot f_2) = f_1 \cdot \mathbf{I}_\mu(f_2) = f_2 \cdot \mathbf{I}_\mu(f_1).$$

From definitions we have also for any $a \in V$ and for any $f \in \mathcal{E}(E)$

$$\mathbf{I}_\mu(T_a(f)) = T_a(\mathbf{I}_\mu(f)).$$

4.4 Fourier transform

Let $E = (I, F, V)$ be a complete C_1 -space over the field \mathbb{F}_q . Then $\check{E} = E$.

Any $a \in V$ defines the function $\psi_a \in \check{\mathcal{E}}(\check{E})$:

$$\psi_a(w) \stackrel{\text{def}}{=} \psi(a(w)), \quad w \in \check{V}.$$

Definition 5 We fix $\mu \in \mu(E)$. We define the Fourier transform $\mathbf{F}_\mu : \mathcal{D}(E) \rightarrow \mathcal{F}(\check{V})$:

$$\mathbf{F}_\mu(f)(w) \stackrel{\text{def}}{=} \int_V f \cdot \overline{\psi_w} d\mu, \quad f \in \mathcal{D}(E), \quad w \in \check{V}.$$

Example 2 Let $i \in I$. Then by easy calculations we obtain that

$$\mathbf{F}_\mu(\delta_{F(i)}) = \mu(F(i)) \cdot \delta_{F(i)^\perp}.$$

Lemma 2 Let $a \in V$, $b \in \check{V}$, $f \in \mathcal{D}(V)$. Then

1.

$$\mathbf{F}_\mu(T_a(f)) = \psi_a \cdot \mathbf{F}_\mu(f),$$

2.

$$\mathbf{F}_\mu(\psi_b \cdot f) = T_{-b} \cdot \mathbf{F}_\mu(f).$$

Proof follows at once from definition of Fourier transform by an easy calculation.

From example 2 and lemma 2 we see that $\mathbf{F}_\mu(T_a(\delta_{F(i)})) \in \mathcal{D}(\check{E})$ for any $a \in V$, $i \in I$. Since the space $\mathcal{D}(E)$ is generated by $T_a(\delta_{F(i)})$, $a \in V$, $i \in I$, we obtain that $\mathbf{F}_\mu(\mathcal{D}(E)) \subset \mathcal{D}(\check{E})$.

The Fourier transform depends linearly on $\mu \in \mu(E)$. Therefore the following map \mathbf{F} is well-defined:

$$\begin{aligned} \mathbf{F} &: \mathcal{D}(E) \otimes_{\mathbf{C}} \mu(E) \longrightarrow \mathcal{D}(\check{E}) \\ \mathbf{F}(f \otimes \mu) &\stackrel{\text{def}}{=} \mathbf{F}_\mu(f), \quad f \in \mathcal{D}(E), \quad \mu \in \mu(E). \end{aligned} \tag{22}$$

For any complete C_1 -space $E = (I, F, V)$ we have a canonical isomorphism

$$\mu(E) \otimes_{\mathbf{C}} \mu(\check{E}) \longrightarrow \mathbf{C}, \tag{23}$$

where if $\mu \in \mu(E)$, $\mu \neq 0$, then $\mu \otimes \mu^{-1} \mapsto 1$, and $\mu^{-1} \in \mu(\check{E}) = \mu(E)^*$, $\mu^{-1}(F(i)^\perp) \stackrel{\text{def}}{=} \mu(F(i))^{-1}$ for any $i \in I$.

Proposition 9 *Let $\mu \in \mu(E)$, $\mu \neq 0$. Then*

1. $\mathbf{F}_{\mu^{-1}} \circ \mathbf{F}_{\mu}(f) = \check{f}$ for any $f \in \mathcal{D}(E)$;
2. \mathbf{F}_{μ} is an isomorphism of \mathbf{C} -vector spaces $\mathcal{D}(E)$ and $\mathcal{D}(\check{E})$.

Proof . The first statement follows by direct calculations on the functions $T_a(\delta_{F(i)})$, $a \in V$, $i \in I$ by using example 2 and lemma 2, because the space $\mathcal{D}(E)$ is generated by the functions $T_a(\delta_{F(i)})$, $a \in V$, $i \in I$.

The second statement of this proposition follows from the first statement.

Definition 6 *We fix $\mu \in \mu(E)$ such that $\mu \neq 0$. We define the Fourier transform $\mathbf{F}_{\mu^{-1}} : \mathcal{D}'(E) \rightarrow \mathcal{D}'(\check{E})$ in the following way:*

$$\mathbf{F}_{\mu^{-1}}(G)(f) \stackrel{\text{def}}{=} G(\mathbf{F}_{\mu^{-1}}(f)), \quad G \in \mathcal{D}'(E), \quad f \in \mathcal{D}(\check{E}).$$

From definitions 5 and 6 we have that

$$\mathbf{F}_{\mu^{-1}} \circ \mathbf{I}_{\mu}(f) = \mathbf{I}_{\mu^{-1}} \circ \mathbf{F}_{\mu}(f), \quad f \in \mathcal{D}(E).$$

From definition 6 and formulas (22) we have that the following map \mathbf{F} is well-defined:

$$\begin{aligned} \mathbf{F} &: \mathcal{D}'(E) \otimes_{\mathbf{C}} \mu(\check{E}) \longrightarrow \mathcal{D}'(\check{E}) \\ \mathbf{F}(G \otimes \mu^{-1}) &\stackrel{\text{def}}{=} \mathbf{F}_{\mu^{-1}}(G), \quad G \in \mathcal{D}(E), \quad \mu \in \mu(E), \quad \mu \neq 0. \end{aligned}$$

Lemma 3 *Let $G \in \mathcal{D}'(E)$, $a \in V$, $b \in \check{V}$. Then*

1.

$$\mathbf{F}_{\mu^{-1}}(T_a(G)) = \psi_a \cdot \mathbf{F}_{\mu^{-1}}(G),$$

2.

$$\mathbf{F}_{\mu^{-1}}(\psi_b \cdot G) = T_{-b}(\mathbf{F}_{\mu^{-1}}(G)).$$

Proof . Using lemma 2, we have for any $f \in \mathcal{D}(\check{E})$:

$$\begin{aligned} \mathbf{F}_{\mu^{-1}}(T_a(G))(f) &= (T_a(G))(\mathbf{F}_{\mu^{-1}}(f)) = G(T_{-a}(\mathbf{F}_{\mu^{-1}}(f))) = \\ &= G(\mathbf{F}_{\mu^{-1}}(\psi_a \cdot f)) = \mathbf{F}_{\mu^{-1}}(G)(\psi_a \cdot f) = (\psi_a \cdot \mathbf{F}_{\mu^{-1}}(G))(f). \end{aligned}$$

$$\begin{aligned} \mathbf{F}_{\mu^{-1}}(\psi_b \cdot G)(f) &= (\psi_b \cdot G)(\mathbf{F}_{\mu^{-1}}(f)) = G(\psi_b \cdot \mathbf{F}_{\mu^{-1}}(f)) = \\ &= G(\mathbf{F}_{\mu^{-1}}(T_b(f))) = \mathbf{F}_{\mu^{-1}}(G)(T_b(f)) = T_{-b}(\mathbf{F}_{\mu^{-1}}(G(f))). \end{aligned}$$

For any $G \in \mathcal{D}'(E)$ we define $\check{G} \in \mathcal{D}'(E)$:

$$\check{G}(f) \stackrel{\text{def}}{=} G(\check{f}), \quad f \in \mathcal{D}(E).$$

We note that for any $g \in \mathcal{E}(E)$ we have

$$\mathbf{I}_{\mu}(\check{g}) = \mathbf{I}_{\mu}(g)^{\check{}}.$$

Proposition 10 *Let $\mu \in \mu(E)$, $\mu \neq 0$. Then*

1. $\mathbf{F}_\mu \circ \mathbf{F}_{\mu^{-1}}(G) = \check{G}$ for any $G \in \mathcal{D}'(E)$;
2. $\mathbf{F}_{\mu^{-1}}$ is an isomorphism of \mathbf{C} -vector spaces $\mathcal{D}'(E)$ and $\mathcal{D}'(\check{E})$.

Proof follows at once from proposition 9.

Now we define the Fourier transform $\mathbf{F} : \mathcal{E}(E) \longrightarrow \mathcal{D}'(\check{E})$ by the following rule:

$$\mathbf{F}(g) \stackrel{\text{def}}{=} \mathbf{F}_{\mu^{-1}} \circ \mathbf{I}_\mu(g), \quad g \in \mathcal{E}(E) . \quad (24)$$

We note that this map \mathbf{F} does not depend on the choice of $\mu \in \mu(E)$, although we use $\mu \in \mu(E)$, $\mu \neq 0$ for the definition of this map \mathbf{F} .

Remark 6 Let $i \leq j \in I$, then in section 3 we defined the Foutier transform

$$\mathbf{F} : \mathcal{F}(F(j)/F(i)) \longrightarrow \mathcal{F}((F(j)/F(i))^*).$$

Now using successively projective and inductive limits from formulas (10)-(15) and explicit description of \check{V} from subsection 4.1, by means of statement 4 of proposition 4 we redefine the Fourier transform $\mathbf{F}_{\mu^{-1}} : \mathcal{D}'(E) \rightarrow \mathcal{D}'(\check{E})$. This new definition of Fourier transform coincides with the definition of Fourier transform which was given in this section above.

Using this definition of Fourier transform, we have the following proposition.

Proposition 11 *The following statements are satisfied.*

1. \mathbf{F} gives an isomorphism of \mathbf{C} -vector spaces $\mathcal{E}(E)$ and $\tilde{\mathcal{E}}'(\check{E})$.
2. $\mathbf{F}|_{\tilde{\mathcal{E}}(E)}$ gives an isomorphism of \mathbf{C} -vector spaces $\tilde{\mathcal{E}}(E)$ and $\mathcal{E}'(\check{E})$.
3. For any $f \in \mathcal{E}(E)$, for any $g \in \tilde{\mathcal{E}}(\check{E})$ we have

$$\langle \mathbf{F}(f), g \rangle_{\tilde{\mathcal{E}}(\check{E})} = \langle f, \mathbf{F}(g) \rangle_{\mathcal{E}(E)} . \quad (25)$$

4. For any $G \in \tilde{\mathcal{E}}'(\check{E})$, for any $v \in V$ we have

$$\mathbf{F}^{-1}(G)(v) = \langle G, \psi_v \rangle_{\tilde{\mathcal{E}}(\check{E})} . \quad (26)$$

Proof . Statements 1, 2 follow from remark 6 and formulas (12)-(15). Statement 3 follows from statement 3 of proposition 4, statements 1-2 of this proposition and remark 4.

Statement 4 follows from statement 3 in the following way. For any $v \in V$ we have

$$\delta_v \in \mathcal{E}'(E) : \quad \delta_v(g) \stackrel{\text{def}}{=} g(v), \quad g \in \mathcal{D}(E).$$

We have $\langle f, \delta_v \rangle_{\mathcal{E}(E)} = f(v)$ for any $f \in \mathcal{E}(E)$. We have also $\mathbf{F}(\psi_v) = T_{-v}(\mathbf{F}(1)) = T_{-v}(\delta_0) = \delta_v$. Now formula (26) follows from formula (25).

4.5 Poisson formula for 1-dimensional case.

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of complete C_1 -spaces over the field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. Then we have the following admissible triple of complete C_1 -spaces:

$$0 \longrightarrow \check{E}_3 \xrightarrow{\check{\beta}} \check{E}_2 \xrightarrow{\check{\alpha}} \check{E}_1 \longrightarrow 0.$$

Besides, we have the following canonical isomorphism:

$$\mu(E_1) \otimes_{\mathbf{C}} \mu(E_3) \longrightarrow \mu(E_2), \quad (27)$$

where $(\mu_1 \otimes \mu_3)(F_2(j)) \stackrel{\text{def}}{=} \mu_1(F_2(j) \cap V_1) \cdot \mu_3(\beta(F_2(j)))$, $j \in I_2$, $\mu_1 \in \mu(V_1)$, $\mu_3 \in \mu(V_3)$.

We define the map α^* :

$$\alpha^* : \mathcal{D}(E_2) \longrightarrow \mathcal{D}(E_1), \quad (28)$$

where $\alpha^*(f)(v) \stackrel{\text{def}}{=} f(\alpha(v))$, $f \in \mathcal{D}(E_2)$, $v \in V_1$. This map is well-defined.

As conjugate map to the map α^* we define the map α_* :

$$\alpha_* : \mathcal{D}'(E_1) \longrightarrow \mathcal{D}'(E_2). \quad (29)$$

We note that if we fixed $\mu_1 \in \mu(E_1)$, $\mu_1 \neq 0$, and fixed $\mu_2 \in \mu(E_2)$, $\mu_2 \neq 0$, then from formulas (23) and (27) we have well-defined $\mu_1 \otimes \mu_2^{-1} \in \mu(\check{E}_3)$.

We define the *characteristic function* of subspace E_1 as distribution $\delta_{E_1, \mu_1} \in \mathcal{D}'(E_2)$, where

$$\delta_{E_1, \mu_1}(f) \stackrel{\text{def}}{=} \int_{V_1} \alpha^*(f) d\mu_1 \quad \text{for any } f \in \mathcal{D}(E_2).$$

Analogously we can define $\delta_{\check{E}_3, \mu_1 \otimes \mu_2^{-1}} \in \mathcal{D}'(\check{E}_2)$.

Theorem 1 (Poisson formula) . For any $f \in \mathcal{D}(E_2)$, for any $\mu_1 \in \mu(E_1)$, $\mu_1 \neq 0$, and $\mu_2 \in \mu(E_2)$, $\mu_2 \neq 0$ the following equivalent statements are satisfied.

1.

$$\int_{V_1} \alpha^*(f) d\mu_1 = \int_{\check{V}_3} \check{\beta}^*(\mathbf{F}_{\mu_2}(f)) d(\mu_1 \otimes \mu_2^{-1}) \quad . \quad (30)$$

2.

$$\mathbf{F}_{\mu_2^{-1}}(\delta_{E_1, \mu_1}) = \delta_{\check{E}_3, \mu_1 \otimes \mu_2^{-1}} \quad . \quad (31)$$

Proof . We have the following obvious properties:

$$T_a(\delta_{E_1, \mu_1}) = \delta_{E_1, \mu_1} \quad \text{for any } a \in V_1 \quad ; \quad (32)$$

$$\psi_b \cdot \delta_{E_1, \mu_1} = \delta_{E_1, \mu_1} \quad \text{for any } b \in \check{V}_3 \quad . \quad (33)$$

There are analogous properties for $\delta_{\tilde{E}_3, \mu_1 \otimes \mu_2^{-1}}$.

By lemma 3 we have that properties given by formulas (32)-(33) are changed with each other under the Fourier transform. We show that δ_{E_1, μ_1} is determined by these properties uniquely up to some constant. (And the analogy is true for $\delta_{\tilde{E}_3, \mu_1 \otimes \mu_2^{-1}}$.) Indeed, from formula (33) it follows that $\delta_{E_1, \mu_1} = \alpha_*(G)$ for some $G \in \mathcal{D}'(E_1)$. Now from formula (32) and proposition 7 we obtain that $G(g) = c \cdot \int_{V_1} g d\mu_1$ for some $c \in \mathbf{C}$ and any $g \in \mathcal{D}(E_1)$.

Therefore we proved formula (31) up to some constant $d \in \mathbf{C}$. To obtain that $d = 1$ we calculate formula (31) on $\delta_{F_2(j)} \in \mathcal{D}(E_2)$ for some $j \in I_2$ using explicit example 2.

4.6 Direct and inverse images.

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_1 -spaces over the field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$.

Now we define the map β_* :

$$\beta_* : \mathcal{D}(E_2) \otimes_{\mathbf{C}} \mu(E_1) \longrightarrow \mathcal{D}(E_3), \quad (34)$$

where

$$\beta_*(f \otimes \mu_1)(w) \stackrel{\text{def}}{=} \int_{\beta^{-1}(w)} f|_{\beta^{-1}(w)} d\mu_1 \stackrel{\text{def}}{=} \int_{V_1} T_{w'}(f) d\mu_1 = \mu_1(T_{w'}(f)|_{V_1}). \quad (35)$$

Here $f \in \mathcal{D}(E_2)$, $\mu_1 \in \mu(E_1)$, $w' \in \beta^{-1}(w)$ and then $\beta^{-1}(w) = w' + V_1$. The result doesn't depend on a choice of the w' .

As conjugate map to the map β_* we define the map β^* :

$$\beta^* : \mathcal{D}'(E_3) \otimes_{\mathbf{C}} \mu(E_1) \longrightarrow \mathcal{D}'(E_2). \quad (36)$$

We have the following property of the map β_* .

Proposition 12 (Fubini formula) *Let $f \in \mathcal{D}(E_2)$, $\mu_i \in \mu(E_i)$ for $1 \leq i \leq 3$. Let $\mu_1 \otimes \mu_3 = \mu_2$ under the map (27). Then*

$$\int_{E_3} \beta_*(f \otimes \mu_1) d\mu_3 = \int_{E_2} f d\mu_2.$$

Proof. We know that the space $\mathcal{D}(E_2)$ is generated by functions $T_a(\delta_{F_2(i)})$, where $a \in V_2$, $i \in I_2$. Therefore we check the Fubini formula on these functions.

Now we extend the direct and inverse images of admissible monomorphisms and epimorphisms (see formulas (28), (29), (34), (36)) to other types of functions and distributions from diagram (21). The problem is that these maps are not defined for arbitrary admissible monomorphisms and epimorphisms.

Proposition 13 . *The following maps of spaces of functions on C_1 -spaces are well-defined.*

1. $\alpha^* : \mathcal{D}(E_2) \longrightarrow \mathcal{D}(E_1).$
2. $\beta^* : \mathcal{D}(E_3) \longrightarrow \mathcal{D}(E_2),$ where E_1 is compact, $V_1 = \text{Ker}(\beta).$
3. $\alpha_* : \mathcal{D}(E_1) \longrightarrow \mathcal{D}(E_2),$ where E_3 is discrete, $V_3 = \text{Coker}(\alpha).$
4. $\beta_* : \mathcal{D}(E_2) \otimes_{\mathbf{C}} \mu(E_1) \longrightarrow \mathcal{D}(E_3).$
5. $\alpha^* : \mathcal{E}(E_2) \longrightarrow \mathcal{E}(E_1),$
and $\alpha^*(\tilde{\mathcal{E}}(E_2)) \subset \tilde{\mathcal{E}}(E_1).$
6. $\beta^* : \mathcal{E}(E_3) \longrightarrow \mathcal{E}(E_2),$
and $\beta^*(\tilde{\mathcal{E}}(E_3)) \subset \tilde{\mathcal{E}}(E_2).$
7. $\alpha_* : \mathcal{E}(E_1) \longrightarrow \mathcal{E}(E_2),$ where E_3 is discrete, $V_3 = \text{Coker}(\alpha),$
and $\alpha_*(\tilde{\mathcal{E}}(E_1)) \subset \tilde{\mathcal{E}}(E_2).$
8. $\beta_* : \mathcal{E}(E_2) \otimes_{\mathbf{C}} \mu(E_1) \longrightarrow \mathcal{E}(E_3),$ where E_1 is compact, $V_1 = \text{Ker}(\beta),$
and $\beta_*(\tilde{\mathcal{E}}(E_2) \otimes_{\mathbf{C}} \mu(E_1)) \subset \tilde{\mathcal{E}}(E_3).$

Proof . For any map $\gamma : V \rightarrow W$ we have the map $\gamma^* : \mathcal{F}(W) \rightarrow \mathcal{F}(V)$ by the rule: $\gamma^*(f)(v) \stackrel{\text{def}}{=} f(\gamma(v)),$ where $f \in \mathcal{F}(W), v \in V.$

For any embedding $\lambda : Y \longrightarrow Z$ we have the map $\lambda_* : \mathcal{F}(Y) \longrightarrow \mathcal{F}(Z)$ by the following rule:

$$\lambda_*(f)(z) = \begin{cases} 0 & , \text{ if } z \notin \lambda(Y) \\ f(y) & , \text{ if } z = \lambda(y), \end{cases}$$

where $f \in \mathcal{F}(Y), z \in Z, y \in Y.$

According to remark 3 we have that $\mathcal{E}(E) = \mathcal{D}(E)$ for any compact C_1 -space $E = (I, F, V).$ Therefore for the compact C_1 -space E_1 it is well defined the map

$$\beta_* : \mathcal{E}(E_2) \otimes_{\mathbf{C}} \mu(E_1) \longrightarrow \mathcal{E}(E_3)$$

by formula (35). Now using proposition 5 we verify the statements of this proposition. The proposition is proved

Now as conjugate maps to the maps described in proposition 13 we define the direct and inverse images of admissible monomorphisms and epimorphisms for the spaces of all distributions $\mathcal{D}'(E_i), 1 \leq i \leq 3$ when these maps are defined. To calculate the corresponding direct and inverse images for the other types of distributions from diagram (21) we have to introduce the topology on the spaces of functions $\mathcal{E}(E)$ and $\tilde{\mathcal{E}}(E)$ for a C_1 -space $E = (I, F, V).$

Let $\{B_l\}$ be a projective system of topological k -vector spaces with surjective transition maps (over a discrete field k). Then $\varprojlim_l B_l$ is a topological k -vector space with a base of neighbourhoods of 0 given by $\phi_l^{-1}(V)$ for all l and all open

k -subspaces V of B_l , where $\phi_l : \lim_{\leftarrow l} B_l \rightarrow B_l$ is the natural map. By $\text{Hom}_k^{cont}(\cdot, \cdot)$ we denote a k -space of all k -linear continuous maps between topological k -vector spaces. Then we have a canonical isomorphism:

$$\text{Hom}_k^{cont}(\lim_{\leftarrow l} B_l, k) = \lim_{\rightarrow l} \text{Hom}_k^{cont}(B_l, k). \quad (37)$$

Let $\{A_m\}$ be an inductive system of topological k -vector spaces with injective transition maps (over a discrete field k). Then $\lim_{\rightarrow m} A_m$ is a topological k -vector space with a base of neighbourhoods of zero given by all k -subspaces $U \subset \lim_{\rightarrow m} A_m$ such that $U \cap A_m$ is an open k -subspace in A_m for any m . Then we have a canonical isomorphism:

$$\text{Hom}_k^{cont}(\lim_{\rightarrow m} A_m, k) = \lim_{\leftarrow m} \text{Hom}_k^{cont}(A_m, k). \quad (38)$$

Now we apply these reasonings to the field $k = \mathbf{C}$ (with a discrete topology) and the \mathbf{C} -vector spaces $\tilde{\mathcal{E}}(E)$ and $\mathcal{E}(E)$. We use formulas (13) and (12). We consider the discrete topology on every finite-dimensional \mathbf{C} -vector space $\mathcal{F}(F(i)/F(j))$ for any $j \leq i \in I$.

Then $\mathcal{E}(E)$ is a topological \mathbf{C} -vector space with the base of neighbourhoods of 0 given by the following \mathbf{C} -subspaces $\mathcal{E}(E)_i$ (for all $i \in I$):

$$\mathcal{E}(E)_i \stackrel{\text{def}}{=} \{f \in \mathcal{E}(E) \text{ such that } f|_{F(i)} = 0\}.$$

Let for any $j \in I$

$$\tilde{\mathcal{E}}(E)_j \stackrel{\text{def}}{=} \{f \in \tilde{\mathcal{E}}(E) \text{ such that } T_w f = f \text{ for any } w \in F(j)\}.$$

Then $\tilde{\mathcal{E}}(E) = \bigcup_{j \in I} \tilde{\mathcal{E}}(E)_j$, and $\tilde{\mathcal{E}}(E)$ is a topological \mathbf{C} -vector space with the base of neighbourhoods of zero given by all \mathbf{C} -subspaces $U \subset \tilde{\mathcal{E}}(E)$ such that for any $j \in I$ there is $i \in I$, $i \geq j$ such that

$$U \cap \tilde{\mathcal{E}}(E)_j \supset \mathcal{E}(E)_i \cap \tilde{\mathcal{E}}(E)_j.$$

We have that the canonical embedding $\tilde{\mathcal{E}}(E) \subset \mathcal{E}(E)$ is a continuous map with respect to these topologies, but the restriction of the topology of $\mathcal{E}(E)$ to $\tilde{\mathcal{E}}(E)$ doesn't coincide with the topology of $\tilde{\mathcal{E}}(E)$.

Using formulas (14) and (15) for the \mathbf{C} -spaces $\mathcal{E}'(E)$ and $\tilde{\mathcal{E}}'(E)$, and also isomorphisms (37) and (38) we obtain the following canonical isomorphisms

$$\mathcal{E}'(E) = \text{Hom}_{\mathbf{C}}^{cont}(\mathcal{E}(E), \mathbf{C}) \quad \text{and} \quad \tilde{\mathcal{E}}'(E) = \text{Hom}_{\mathbf{C}}^{cont}(\tilde{\mathcal{E}}(E), \mathbf{C}). \quad (39)$$

We have the following lemma.

Lemma 4 *Suppose that conditions of proposition 13 are satisfied. Then*

1. The maps α_* , α^* , β_* , β^* (when they are defined) are continuous on the topological \mathbf{C} -vector spaces $\mathcal{E}(E_i)$, $1 \leq i \leq 3$.
2. The maps α_* , α^* , β_* , β^* (when they are defined) are continuous on the topological \mathbf{C} -vector spaces $\tilde{\mathcal{E}}(E_i)$, $1 \leq i \leq 3$.

Proof . We use explicit descriptions of open subsets on $\mathcal{E}(E)$ and $\tilde{\mathcal{E}}(E)$ given above. For example, to prove that β_* is a continuous map from \mathbf{C} -vector space $\mathcal{E}(E_2)$ to \mathbf{C} -vector space $\mathcal{E}(E_3)$ (or from \mathbf{C} -vector space $\tilde{\mathcal{E}}(E_2)$ to \mathbf{C} -vector space $\tilde{\mathcal{E}}(E_3)$) when E_1 is a compact C_1 -space, we use that

$$\beta_*(\mathcal{E}(E_2)_i \otimes_{\mathbf{C}} \mu(E_1)) \subset \mathcal{E}(E_3)_{i'}$$

when $i \in I_2$, $F_2(i) \supset V_1$, and $i' \in I_3$, $F_3(i') = \beta(F_2(i))$. And also

$$\beta_*(\tilde{\mathcal{E}}(E_2)_j \otimes_{\mathbf{C}} \mu(E_1)) \subset \tilde{\mathcal{E}}(E_3)_{j'}$$

when $j \in I_2$, $j' \in I_3$, and $F_3(j') = \beta(F_2(j))$.

Using this lemma and formulas (39) we consider the direct and inverse images as conjugate maps to the maps considered in proposition 13.

Proposition 14 . The following maps of spaces of distributions on C_1 -spaces are well-defined.

1. α_* : $\mathcal{D}'(E_1) \longrightarrow \mathcal{D}'(E_2)$.
2. β_* : $\mathcal{D}'(E_2) \longrightarrow \mathcal{D}'(E_3)$, where E_1 is compact, $V_1 = \text{Ker}(\beta)$.
3. α^* : $\mathcal{D}'(E_2) \longrightarrow \mathcal{D}'(E_1)$, where E_3 is discrete, $V_3 = \text{Coker}(\alpha)$.
4. β^* : $\mathcal{D}'(E_3) \otimes_{\mathbf{C}} \mu(E_1) \longrightarrow \mathcal{D}'(E_2)$.
5. α_* : $\tilde{\mathcal{E}}'(E_1) \longrightarrow \tilde{\mathcal{E}}'(E_2)$,
and $\alpha_*(\mathcal{E}'(E_1)) \subset \mathcal{E}'(E_2)$.
6. β_* : $\tilde{\mathcal{E}}'(E_2) \longrightarrow \tilde{\mathcal{E}}'(E_3)$,
and $\beta_*(\mathcal{E}'(E_2)) \subset \mathcal{E}'(E_3)$.
7. α^* : $\tilde{\mathcal{E}}'(E_2) \longrightarrow \tilde{\mathcal{E}}'(E_1)$, where E_3 is discrete, $V_3 = \text{Coker}(\alpha)$,
and $\alpha^*(\mathcal{E}'(E_2)) \subset \mathcal{E}'(E_1)$.
8. β^* : $\tilde{\mathcal{E}}'(E_3) \otimes_{\mathbf{C}} \mu(E_1) \longrightarrow \tilde{\mathcal{E}}'(E_2)$, where E_1 is compact, $V_1 = \text{Ker}(\beta)$,
and $\beta^*(\mathcal{E}'(E_3) \otimes_{\mathbf{C}} \mu(E_1)) \subset \mathcal{E}'(E_2)$.

Proof . We use proposition 13, lemma 4, and formulas (39).

Remark 7 Using remark 4 we obtain that conjugate maps defined in proposition 14 are compatible with respect to the canonical embeddings:

$$\mathcal{E}'(E_i) \subset \tilde{\mathcal{E}}'(E_i) \subset \mathcal{D}'(E_i), \quad 1 \leq i \leq 3.$$

From definitions of the maps α^* , α_* , β^* , β_* we have the following *projections formulas*. We fix $\mu_1 \in \mu(E_1)$.

For any $f \in \mathcal{D}(E_2)$ and $g \in \mathcal{E}(E_3)$

$$\beta_*(f \cdot \beta^*(g) \otimes \mu_1) = \beta_*(f \otimes \mu_1) \cdot g. \quad (40)$$

For any $f \in \mathcal{E}(E_2)$, $g \in \mathcal{E}(E_3)$, and a compact C_1 -space E_1 we have

$$\beta_*(f \cdot \beta^*(g) \otimes \mu_1) = \beta_*(f \otimes \mu_1) \cdot g. \quad (41)$$

For any $f \in \mathcal{E}(E_1)$, $g \in \mathcal{E}(E_2)$, and a discrete C_1 -space E_3

$$\alpha_*(f \cdot \alpha^*(g)) = \alpha_*(f) \cdot g. \quad (42)$$

Proposition 15 *Let $\mu_i \in \mu(E_i)$, $1 \leq i \leq 3$ such that $\mu_1 \otimes \mu_3 = \mu_2$ according to map (27). We have the following formulas.*

1. $\mathbf{I}_{\mu_2}(\beta^*(g)) = \beta^*(\mathbf{I}_{\mu_3}(g))$ for any $g \in \mathcal{E}(E_3)$.
2. Let E_1 be a compact C_1 -space. Then $\mathbf{I}_{\mu_3}(\beta_*(f \otimes \mu_1)) = \beta_*(\mathbf{I}_{\mu_2}(f))$ for any $f \in \mathcal{E}(E_2)$.

Proof follows from Fubini formula (proposition 12) and projection formulas (40) and (41).

4.7 Composition of maps and base change rules.

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_1 -spaces over the field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$.

Let

$$0 \longrightarrow L \xrightarrow{\alpha'} H \xrightarrow{\beta'} E_2 \longrightarrow 0$$

be another admissible triple of C_1 -spaces over the field \mathbb{F}_q , where $L = (J_1, T_1, U)$, $H = (J_2, T_2, W)$.

We have then the following admissible triple of C_1 -spaces:

$$0 \longrightarrow H \times_{E_2} E_1 \xrightarrow{\beta'_\alpha} H \xrightarrow{\beta\beta'} E_3 \longrightarrow 0.$$

Let $H' = H \times_{E_2} E_1 = (K, P, Z)$ as C_1 -space.

Proposition 16 . *We have the following formulas.*

1. For any $f \in \mathcal{D}(H)$, $\nu \in \mu(L)$, $\mu \in \mu(E_1)$

$$(\beta\beta')_*(f \otimes (\nu \otimes \mu)) = \beta_*(\beta'_*(f \otimes \nu) \otimes \mu). \quad (43)$$

2. For any $G \in \mathcal{D}'(E_3)$, $\nu \in \mu(L)$, $\mu \in \mu(E_1)$

$$(\beta\beta')^*(G \otimes (\nu \otimes \mu)) = (\beta')^*(\beta^*(G \otimes \mu) \otimes \nu). \quad (44)$$

3. For any $f \in \mathcal{E}(E_3)$

$$(\beta\beta')^*(f) = (\beta')^*\beta^*(f). \quad (45)$$

4. For any $G \in \tilde{\mathcal{E}}'(H)$

$$(\beta\beta')_*(G) = \beta_*\beta'_*(G). \quad (46)$$

We suppose further that E_1 and L are compact C_1 -spaces. Then $H \times_{E_2} E_1$ is a compact C_1 -space, and the following formulas are satisfied.

5. For any $f \in \mathcal{D}(E_3)$

$$(\beta\beta')^*(f) = (\beta')^*\beta^*(f). \quad (47)$$

6. For any $G \in \mathcal{D}'(H)$

$$(\beta\beta')_*(G) = \beta_*(\beta')_*(G). \quad (48)$$

7. For any $f \in \mathcal{E}(H)$, $\nu \in \mu(L)$, $\mu \in \mu(E_1)$

$$(\beta\beta')_*(f \otimes (\nu \otimes \mu)) = \beta_*(\beta'_*(f \otimes \nu) \otimes \mu). \quad (49)$$

8. For any $G \in \tilde{\mathcal{E}}'(E_3)$, $\nu \in \mu(L)$, $\mu \in \mu(E_1)$

$$(\beta\beta')^*(G \otimes (\nu \otimes \mu)) = (\beta')^*(\beta^*(G \otimes \mu) \otimes \nu). \quad (50)$$

Proof . We have by formula (27) that the following equalities are canonically satisfied:

$$\begin{aligned} \mu(E_1) \otimes_{\mathbf{C}} \mu(E_3) &= \mu(E_2) \quad , \quad \mu(L) \otimes_{\mathbf{C}} \mu(E_2) = \mu(H), \\ \mu(H') \otimes_{\mathbf{C}} \mu(E_3) &= \mu(H). \end{aligned}$$

Therefore we have that

$$\mu(L) \otimes_{\mathbf{C}} \mu(E_1) = \mu(H').$$

The last formula follows also from the following admissible triple of C_1 -spaces:

$$0 \longrightarrow L \xrightarrow{\alpha_{\alpha'}} H \times_{E_2} E_1 \xrightarrow{\alpha_{\beta'}} E_1 \longrightarrow 0. \quad (51)$$

Therefore for $\nu \in \mu(L)$, $\mu \in \mu(E_1)$ we have canonically $\nu \otimes \mu \in \mu\left(H \times_{E_2} E_1\right)$.

Now formula (43) follows from Fubini formula (proposition 12). Formula (44) is the conjugate formula to formula (43). Formula (45) is evident. We restrict formula (45) to the subspace $\tilde{\mathcal{E}}(E_3)$. Then formula (46) is the conjugate formula to the last formula.

From triple (51) we obtain that if E_1 and L are compact C_1 -spaces, then $H \times_{E_2} E_1$ is also a compact C_1 -space.

Now formula (47) follows from the definitions of inverse images. Formula (48) is the conjugate formula to formula (47). Formula (49) follows from Fubini formula (proposition 12). Formula (50) follows from formula (44). The proof is finished.

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_1 -spaces over the field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. Let $\alpha' : E_2 \longrightarrow H'$ be an admissible monomorphism, where $H' = (J', T', W')$, that is, there is an admissible triple of C_1 -spaces

$$0 \longrightarrow E_2 \xrightarrow{\alpha'} H' \xrightarrow{\beta'} L' \longrightarrow 0.$$

We have the following admissible triple of C_1 -spaces

$$0 \longrightarrow E_1 \xrightarrow{\alpha' \alpha} H' \xrightarrow{\alpha' \beta} E_3 \amalg_{E_2} H' \longrightarrow 0.$$

Proposition 17 . *We have the following formulas.*

1. For any $f \in \mathcal{D}(H')$

$$(\alpha' \alpha)^*(f) = \alpha^*(\alpha')^*(f). \quad (52)$$

2. For any $G \in \mathcal{D}'(E_1)$

$$(\alpha' \alpha)_*(G) = (\alpha')_* \alpha_*(G). \quad (53)$$

3. For any $f \in \mathcal{E}(H')$

$$(\alpha' \alpha)^*(f) = \alpha^*(\alpha')^*(f). \quad (54)$$

4. For any $G \in \tilde{\mathcal{E}}'(E_1)$

$$(\alpha' \alpha)_*(G) = (\alpha')_* \alpha_*(G). \quad (55)$$

We suppose further that E_3 and L' are discrete C_1 -spaces. Then $E_3 \amalg_{E_2} H'$ is a discrete C_1 -space, and the following formulas are satisfied:

5. For any $f \in \mathcal{D}(E_1)$

$$(\alpha' \alpha)_*(f) = (\alpha')_* \alpha_*(f). \quad (56)$$

6. For any $G \in \mathcal{D}'(H')$

$$(\alpha' \alpha)^*(G) = \alpha^*(\alpha')^*(G). \quad (57)$$

7. For any $f \in \mathcal{E}(E_1)$

$$(\alpha' \alpha)_*(f) = (\alpha')_* \alpha_*(f). \quad (58)$$

8. For any $G \in \tilde{\mathcal{E}}'(H')$

$$(\alpha' \alpha)^*(G) = \alpha^*(\alpha')^*(G). \quad (59)$$

Proof . Formula (52) follows from definitions of the corresponding maps. Formula (53) is the conjugate formula to formula (52). Formula (54) follows also from definitions of the corresponding maps. We restrict formula (54) to the subspace $\tilde{\mathcal{E}}(W')$. Then formula (55) is the conjugate formula to this last formula.

From admissible triple of C_1 -spaces

$$0 \longrightarrow E_3 \xrightarrow{\beta_{\alpha'}} E_3 \amalg_{E_2} H' \xrightarrow{\beta_{\beta'}} L' \longrightarrow 0$$

we obtain that if E_3 and L' are discrete C_1 -spaces, then $E_3 \amalg_{E_2} H'$ is also a discrete C_1 -space.

Now formulas (56) and (58) follow at once from definitions. Formula (57) is the conjugate formula to formula (56). Formula (59) follows from formula (57). The proposition is proved.

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_1 -spaces over the field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. Let $\gamma : D \longrightarrow E_3$ be an admissible monomorphism, where $D = (R, S, Y)$. That is, there is the following admissible triple of C_1 -spaces

$$0 \longrightarrow D \xrightarrow{\gamma} E_3 \xrightarrow{\delta} B \longrightarrow 0.$$

Then we have the following commutative diagram (see section 4.1):

$$\begin{array}{ccc} E_2 \times_{E_3} D & \xrightarrow{\gamma_\beta} & D \\ \beta_\gamma \downarrow & & \downarrow \gamma \\ E_2 & \xrightarrow{\beta} & E_3 \end{array}$$

where γ_β is an admissible epimorphism, and β_γ is an admissible monomorphism. Let $X' = E_2 \times_{E_3} D = (N, Q, X)$ as a C_1 -space.

Proposition 18 (*Base change rules*) . We have the following formulas.

1. For any $f \in \mathcal{D}(E_2)$, $\mu \in \mu(E_1)$

$$\gamma^* \beta_*(f \otimes \mu) = (\gamma_\beta)_*(\beta_\gamma^*(f) \otimes \mu). \quad (60)$$

2. For any $G \in \mathcal{D}'(D)$, $\mu \in \mu(E_1)$

$$\beta^*(\gamma_*(G) \otimes \mu) = (\beta_\gamma)_* \gamma_\beta^*(G \otimes \mu). \quad (61)$$

3. For any $f' \in \mathcal{E}(E_3)$

$$\gamma_\beta^* \gamma^*(f') = \beta_\gamma^* \beta^*(f'). \quad (62)$$

4. For any $G' \in \tilde{\mathcal{E}}'(X')$

$$\beta_*(\beta_\gamma)_*(G') = \gamma_*(\gamma_\beta)_*(G'). \quad (63)$$

5. For any $f' \in \mathcal{D}(E_3)$ and compact C_1 -space E_1

$$\gamma_\beta^* \gamma^*(f') = \beta_\gamma^* \beta^*(f'). \quad (64)$$

6. For any $G' \in \mathcal{D}'(X')$ and compact C_1 -space E_1

$$\beta_*(\beta_\gamma)_*(G') = \gamma_*(\gamma_\beta)_*(G'). \quad (65)$$

7. For any $f \in \mathcal{D}(X')$, $\mu \in \mu(E_1)$ and discrete C_1 -space B

$$\beta_*((\beta_\gamma)_*(f) \otimes \mu) = \gamma_*(\gamma_\beta)_*(f \otimes \mu). \quad (66)$$

8. For any $G \in \mathcal{D}'(E_3)$, $\mu \in \mu(E_1)$ and discrete C_1 -space B

$$\gamma_\beta^*(\gamma^*(G) \otimes \mu) = \beta_\gamma^*\beta^*(G \otimes \mu). \quad (67)$$

We suppose further that E_1 is a compact C_1 -space, and B is a discrete C_1 -space. Then the following formulas are satisfied.

9. For any $f \in \mathcal{D}(D)$

$$\beta^*\gamma_*(f) = (\beta_\gamma)_*\gamma_\beta^*(f). \quad (68)$$

10. For any $G \in \mathcal{D}'(E_2)$

$$\gamma^*\beta_*(G) = (\gamma_\beta)_*\beta_\gamma^*(G). \quad (69)$$

11. For any $f \in \mathcal{E}(X')$, $\mu \in \mu(E_1)$

$$\beta_*((\beta_\gamma)_*(f) \otimes \mu) = \gamma_*(\gamma_\beta)_*(f \otimes \mu). \quad (70)$$

12. For any $G \in \tilde{\mathcal{E}}'(E_3)$, $\mu \in \mu(E_1)$

$$\gamma_\beta^*(\gamma^*(G) \otimes \mu) = \beta_\gamma^*\beta^*(G \otimes \mu). \quad (71)$$

Proof . Formula (60) follows from the corresponding definitions. Formula (61) is the conjugate formula to formula (60). Formula (62) follows also from the corresponding definitions. To obtain formula (63), we restrict formula (62) to the subspace $\tilde{\mathcal{E}}(E_3)$ and then take the conjugate formula to this formula. Formulas (64) and (66) follow at once from the corresponding definitions. Formula (65) is the conjugate formula to formula (64), formula (67) is the conjugate formula to formula (66). Formulas (68) and (70) follow at once from the corresponding definitions. Formula (69) is the conjugate formula to formula (68). Formula (71) follows from formula (67).

4.8 Direct and inverse images and Fourier transform.

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of complete C_1 -spaces over the field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. Then we have the following admissible triple of complete C_1 -spaces:

$$0 \longrightarrow \check{E}_3 \xrightarrow{\check{\beta}} \check{E}_2 \xrightarrow{\check{\alpha}} \check{E}_1 \longrightarrow 0.$$

Proposition 19 *We have the following commutative diagrams.²:*

²Here and in the sequel, the expressions of type $\beta_* \otimes \mu(E_3)$ are shortenings of more correct $\beta_* \otimes \text{Id}_{\mu(E_3)}$

$$\begin{array}{ccc}
\mathcal{D}(E_2) \otimes_{\mathbf{C}} \mu(E_2) & \xrightarrow{\beta_* \otimes \mu(E_3)} & \mathcal{D}(E_3) \otimes_{\mathbf{C}} \mu(E_3) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F}
\end{array} \tag{72}$$

$$\begin{array}{ccc}
\mathcal{D}(\check{E}_2) & \xrightarrow{\check{\beta}^*} & \mathcal{D}(\check{E}_3) \\
\mathcal{D}(E_2) & \xrightarrow{\alpha^*} & \mathcal{D}(E_1) \\
\mu(\check{E}_2) \otimes \mathbf{F} \downarrow & & \downarrow \mathbf{F} \otimes \mu(\check{E}_1)
\end{array} \tag{73}$$

$$\begin{array}{ccc}
\mathcal{D}(\check{E}_2) \otimes_{\mathbf{C}} \mu(\check{E}_2) & \xrightarrow{\check{\alpha}_* \otimes \mu(\check{E}_1)} & \mathcal{D}(\check{E}_1) \otimes_{\mathbf{C}} \mu(\check{E}_1) \\
\mathcal{D}'(E_3) \otimes_{\mathbf{C}} \mu(E_3)^* & \xrightarrow{\beta^* \otimes \mu(E_2)^*} & \mathcal{D}'(E_2) \otimes_{\mathbf{C}} \mu(E_2)^* \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F}
\end{array} \tag{74}$$

$$\begin{array}{ccc}
\mathcal{D}'(\check{E}_3) & \xrightarrow{\check{\beta}_*} & \mathcal{D}'(\check{E}_2) \\
\mathcal{D}'(E_1) & \xrightarrow{\alpha_*} & \mathcal{D}'(E_2) \\
\mu(E_1) \otimes \mathbf{F} \downarrow & & \downarrow \mathbf{F} \otimes \mu(E_2)
\end{array} \tag{75}$$

$$\mathcal{D}'(\check{E}_1) \otimes_{\mathbf{C}} \mu(E_1) \xrightarrow{\check{\alpha}_* \otimes \mu(E_1)} \mathcal{D}'(\check{E}_2) \otimes_{\mathbf{C}} \mu(E_2)$$

Proof . We use formulas which are analogous to formulas (23) and (27):

$$\begin{aligned}
\mu(E_1) \otimes_{\mathbf{C}} \mu(E_3) &= \mu(E_2), \\
\mu(\check{E}_3) \otimes_{\mathbf{C}} \mu(\check{E}_1) &= \mu(\check{E}_2), \\
\mu(E_i) \otimes_{\mathbf{C}} \mu(\check{E}_i) &= \mathbf{C}, \quad 1 \leq i \leq 3.
\end{aligned}$$

Now diagram (72) is equivalent to the following formula:

$$\int_{V_3} \beta_*(f) \cdot \overline{\psi_w} d\mu_3 = \int_{V_2} f \cdot \overline{\psi_{\check{\beta}(w)}} d\mu_2, \tag{76}$$

which has to be satisfied for any $f \in \mathcal{D}(E_2)$, any $w \in \check{V}_3$, and $\mu_3 = \mu_1 \otimes \mu_2$.

Now using $\overline{\psi_{\check{\beta}(w)}} = \beta^*(\overline{\psi_w})$, projection formula (40) and Fubini formula (proposition 12), we obtain formula (76).

Diagram (73) follows from diagram (72) and proposition 9. Diagram (74) is the conjugate diagram to diagram (73). Diagram (75) is the conjugate diagram to diagram (72).

Proposition 20 *We have the following commutative diagrams.*

$$\begin{array}{ccc}
\mathcal{E}(E_3) & \xrightarrow{\beta^*} & \mathcal{E}(E_2) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F} \\
\tilde{\mathcal{E}}'(\check{E}_3) & \xrightarrow{\check{\beta}_*} & \tilde{\mathcal{E}}'(\check{E}_2)
\end{array} \tag{77}$$

$$\begin{array}{ccc}
\tilde{\mathcal{E}}'(E_1) & \xrightarrow{\alpha_*} & \tilde{\mathcal{E}}'(E_2) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F} \\
\mathcal{E}(\check{E}_1) & \xrightarrow{\check{\alpha}^*} & \mathcal{E}(\check{E}_2)
\end{array} \tag{78}$$

$$\begin{array}{ccc}
\mathcal{E}(E_2) & \xrightarrow{\alpha^*} & \mathcal{E}(E_1) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F} \\
\tilde{\mathcal{E}}'(\check{E}_2) & \xrightarrow{\check{\alpha}_*} & \tilde{\mathcal{E}}'(\check{E}_1)
\end{array} \tag{79}$$

$$\begin{array}{ccc}
\tilde{\mathcal{E}}'(E_2) & \xrightarrow{\beta_*} & \tilde{\mathcal{E}}'(E_3) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F} \\
\mathcal{E}(\check{E}_2) & \xrightarrow{\check{\beta}^*} & \mathcal{E}(\check{E}_3)
\end{array} \tag{80}$$

Proof . Diagram (77) follows from diagram (74), proposition 15, formula (24), and proposition 11.

Diagram (78) follows from diagram (77), proposition 11, and the following formula: $\mathbf{F} \circ \mathbf{F}(g) = \check{g}$ for any $g \in \mathcal{E}(E_i)$ (which follows from proposition 10).

We change in diagram (77) the spaces $\mathcal{E}(E_i)$ on their subspaces $\tilde{\mathcal{E}}(E_i)$ ($i = 2, 3$) correspondingly, and the spaces $\tilde{\mathcal{E}}'(\check{E}_i)$ also on their subspaces $\mathcal{E}'(\check{E}_i)$ ($i = 2, 3$) correspondingly. Then by proposition 11, this new diagram is commutative. And the conjugation to this diagram gives diagram 79. (We used remark 4.)

Diagram (80) follows from diagram (79) and the following formula: $\mathbf{F} \circ \mathbf{F}(g) = \check{g}$ for any $g \in \mathcal{E}(E_i)$.

4.9 The invariance property.

For any C_1 -space $E = (I, F, V)$ we defined the spaces $\mu(E)$, $\mathcal{D}(E)$, $\mathcal{D}'(E)$, $\mathcal{E}(E)$, $\mathcal{E}'(E)$, $\tilde{\mathcal{E}}(E)$, $\tilde{\mathcal{E}}'(E)$.

Proposition 21 . *Let E' be another C_1 -space such that $E \sim E'$. Then there are the following canonical isomorphisms: $\mu(E) = \mu(E')$,*

$$\begin{aligned} \mathcal{D}(E) &= \mathcal{D}(E') \quad , \quad \mathcal{D}'(E) = \mathcal{D}'(E') \quad , \quad \mathcal{E}(E) = \mathcal{E}(E'), \\ \mathcal{E}'(E) &= \mathcal{E}'(E') \quad , \quad \tilde{\mathcal{E}}(E) = \tilde{\mathcal{E}}(E') \quad , \quad \tilde{\mathcal{E}}'(E) = \tilde{\mathcal{E}}'(E'). \end{aligned}$$

Proof follows from the corresponding definitions.

Remark 8 . The isomorphisms from proposition 21 are well-defined with respect to isomorphisms (23) and (27), direct and inverse images, and Fourier transform.

5 Two-dimensional case

5.1 C_2 -spaces

We recall the definition of C_2 -space from [13].

We fix a field k . By definition, objects of the category C_2 , i.e. $\text{Ob}(C_2)$, are filtered k -vector spaces (I, F, V) with the following additional structures

- (i) for any $i \leq j \in I$ on the k -vector space $F(j)/F(i)$ it is given a structure $E_{i,j} \in \text{Ob}(C_1)$,
- (ii) for any $i \leq j \leq k \in I$

$$0 \longrightarrow E_{i,j} \longrightarrow E_{i,k} \longrightarrow E_{j,k} \longrightarrow 0$$

is an admissible triple from C_1 .

Let $E_1 = (I_1, F_1, V_1)$ and $E_2 = (I_2, F_2, V_2)$ be from $\text{Ob}(C_2)$. Then, by definition, $\text{Mor}_{C_2}(E_1, E_2)$ consists of elements $A \in \text{Hom}_k(V_1, V_2)$ such that the following conditions hold:

- (i) for any $i \in I_1$ there is an $j \in I_2$ such that $A(F_1(i)) \subset F_2(j)$,
- (ii) for any $j \in I_2$ there is an $i \in I_1$ such that $A(F_1(i)) \subset F_2(j)$,
- (iii) for any $i_1 \leq i_2 \in I_1$ and $j_1 \leq j_2 \in I_2$ such that $A(F_1(i_1)) \subset F_2(j_1)$ and $A(F_1(i_2)) \subset F_2(j_2)$ we have that the induced k -linear map

$$\bar{A} : \frac{F_1(i_2)}{F_1(i_1)} \longrightarrow \frac{F_2(j_2)}{F_2(j_1)}$$

is an element from

$$\text{Mor}_{C_1}\left(\frac{F_1(i_2)}{F_1(i_1)}, \frac{F_2(j_2)}{F_2(j_1)}\right).$$

Let $E_1 = (I_1, F_1, V_1)$, $E_2 = (I_2, F_2, V_2)$ and $E_3 = (I_3, F_3, V_3)$ be from $\text{Ob}(C_2)$. Then we say that

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0 \quad (81)$$

is an *admissible triple* from C_2 when the following conditions are satisfied:

(i)

$$0 \longrightarrow V_1 \xrightarrow{\alpha} V_2 \xrightarrow{\beta} V_3 \longrightarrow 0$$

is an exact triple of k -vector spaces,

(ii) the filtration (I_1, F_1, V_1) dominates the filtration (I_2, F'_1, V_1) , where $F'_1(i) = F_2(i) \cap V_1$ for any $i \in I_2$,

(iii) the filtration (I_3, F_3, V_3) dominates the filtration (I_2, F'_3, V_3) , where $F'_3(i) = F_2(i)/F_2(i) \cap V_1$ for any $i \in I_2$,

(iv) for any $i \leq j \in I_2$

$$0 \longrightarrow \frac{F'_1(j)}{F'_1(i)} \longrightarrow \frac{F_2(j)}{F_2(i)} \longrightarrow \frac{F'_3(j)}{F'_3(i)} \longrightarrow 0 \quad (82)$$

is an admissible triple from C_1 . (By definition of $\text{Ob}(C_2)$, on every vector space from triple (82) it is given the structure of $\text{Ob}(C_1)$).

We say that α from an admissible triple (81) is an *admissible monomorphism*, and β from an admissible triple (81) is an *admissible epimorphism*.

Remark 9 Let $E_2 = (I_2, F_2, V_2)$ be a C_2 -space over the field k . Let

$$0 \longrightarrow V_1 \xrightarrow{\alpha} V_2 \xrightarrow{\beta} V_3 \longrightarrow 0 \quad (83)$$

be an exact triple of k -vector spaces. We define on spaces V_1 and V_3 the structures of filtered spaces (I_i, F_i, V_i) , $i = 1, 3$, where $I_1 = I_3 = I_2$, and $F_1(i) \stackrel{\text{def}}{=} F_2(i) \cap V_1$, $F_3(i) \stackrel{\text{def}}{=} \beta(F_2(i))$ for $i \in I_2$.

By definition, for any $i \leq j \in I_2$ on the space $F_2(j)/F_2(i)$ it is given the structure of C_1 -space, and we have the following exact triple of k -vector spaces:

$$0 \longrightarrow F_1(j)/F_1(i) \longrightarrow F_2(j)/F_2(i) \longrightarrow F_3(j)/F_3(i) \longrightarrow 0. \quad (84)$$

Therefore, as in the case of remark 2, we consider the induced filtration on the k -space $F_1(j)/F_1(i)$ and the factor-filtration on the k -space $F_3(j)/F_3(i)$.

The filtrations, so defined, gives always the structure of C_2 -space $E_1 = (I_1, F_1, V_1)$. At the same time, the factor-filtrations, which are defined above, will give the structure of C_2 -space $E_3 = (I_3, F_3, V_3)$ iff the following conditions hold:

(**) for any $x \in V_2 \setminus V_1$ there exist $l \in I_2$ such that $(x + F_2(l)) \cap V_1 = \emptyset$;

(***) for any $i \leq j \in I_2$ for the exact triple (84)

condition (*) from remark 2 is satisfied.

Moreover, under conditions (**), (***), exact triple (83) gives the admissible triple

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0,$$

i.e. α will be an admissible monomorphism, β will be an admissible epimorphism.

We say that a C_2 -space $E_1 = (I_1, F_1, V)$ *dominates* another C_2 -space $E_2 = (I_2, F_2, V)$ if the following conditions are satisfied:

- (i) the filtration (I_1, F_1, V) dominates the filtration (I_2, F_2, V) ,
- (ii) for any $i \leq j \in I_2$ the C_1 -space $E_{1i,j} = F_2(j)/F_2(i)$ with filtration induced by C_2 -structure on E_1 dominates the C_1 -space $E_{2i,j} = F_2(j)/F_2(i)$ with filtration induced by C_2 -structure on E_2 .

We consider the group $\text{Aut}_{C_2}(E) \stackrel{\text{def}}{=} \text{Mor}_{C_2}(E, E)^*$ (i.e., invertible elements in k -algebra $\text{Mor}_{C_2}(E, E)$) for any C_2 -space E . By [13, prop.2.1], if a C_2 -space E_1 dominates a C_1 -space E_2 , then canonically $\text{Aut}_{C_2}(E_1) = \text{Aut}_{C_2}(E_2)$.

Let $E = (I, F, V)$ be a C_2 -space. We define a C_2 -space $\check{E} = (I^0, F^0, \check{V})$ by the following way. The space $\check{V} \subset V^*$ is defined as

$$\check{V} \stackrel{\text{def}}{=} \lim_{\substack{\rightarrow \\ j \in I}} \lim_{\substack{\leftarrow \\ i \geq j}} \check{E}_{j,i}, \quad (85)$$

where the C_1 -space $\check{E}_{j,i}$ is constructed from the C_1 -space $E_{j,i} = F(i)/F(j)$ (see section 4.1). The set I^0 is a partially ordered set, which has the same set as I , but with the inverse order than I . For $j \in I^0$ the subspace

$$F^0(j) \stackrel{\text{def}}{=} \lim_{\substack{\leftarrow \\ i \leq j \in I^0}} \check{E}_{j,i} \subset \check{V}. \quad (86)$$

If $E_1, E_2 \in \text{Ob}(C_2)$ and $\theta \in \text{Mor}_{C_2}(E_1, E_2)$, then there is canonically $\check{\theta} \in \text{Mor}_{C_2}(\check{E}_2, \check{E}_1)$. If

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

is an admissible triple of C_2 -spaces, then there is canonically the following admissible triple of C_2 -spaces:

$$0 \longrightarrow \check{E}_3 \xrightarrow{\check{\beta}} \check{E}_2 \xrightarrow{\check{\alpha}} \check{E}_1 \longrightarrow 0.$$

Lemma 5 *Let*

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_2 -spaces.

1. *Let $D \in \text{Ob}(C_2)$, and $\gamma \in \text{Mor}_{C_2}(D, E_3)$. Then there is the following admissible triple of C_2 -spaces*

$$0 \longrightarrow E_1 \xrightarrow{\gamma_\alpha} E_2 \times_{E_3} D \xrightarrow{\gamma_\beta} D \longrightarrow 0 \quad (87)$$

and $\beta_\gamma \in \text{Mor}_{C_2}(E_2 \times_{E_3} D, E_2)$ such that the following diagram is commutative:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & E_1 & \xrightarrow{\gamma_\alpha} & E_2 \times_{E_3} D & \xrightarrow{\gamma_\beta} & D & \longrightarrow & 0 \\ & & \parallel & & \downarrow \beta_\gamma & & \downarrow \gamma & & \\ 0 & \longrightarrow & E_1 & \xrightarrow{\alpha} & E_2 & \xrightarrow{\beta} & E_3 & \longrightarrow & 0 \end{array} \quad (88)$$

2. Let $A \in \text{Ob}(C_2)$, and $\theta \in \text{Mor}_{C_2}(E_1, A)$. Then there is the following admissible triple of C_2 -spaces

$$0 \longrightarrow A \xrightarrow{\theta_\alpha} A \amalg_{E_1} E_2 \xrightarrow{\theta_\beta} E_3 \longrightarrow 0 \quad (89)$$

and $\alpha_\theta \in \text{Mor}_{C_2}(E_2, A \amalg_{E_1} E_2)$ such that the following diagram is commutative:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & E_1 & \xrightarrow{\alpha} & E_2 & \xrightarrow{\beta} & E_3 & \longrightarrow & 0 \\ & & \downarrow \theta & & \downarrow \alpha_\theta & & \parallel & & \\ 0 & \longrightarrow & A & \xrightarrow{\theta_\alpha} & A \amalg_{E_1} E_2 & \xrightarrow{\theta_\beta} & E_3 & \longrightarrow & 0 \end{array} \quad (90)$$

Proof follows by repeating of reasonings of the proof of lemma 1 when in this proof the fibered product (or the sum) of factors of filtration is understood as the fibered product (or the sum) of C_1 -spaces, constructed in lemma 1.

Definition 7 We say that a C_2 -space (I_1, F_1, V_1) is a cC_2 -space iff there is an $i \in I_1$ such that $F_1(i) = V_1$.

We say that a C_2 -space (I_2, F_2, V_2) is an dC_2 -space iff there is an $j \in I_2$ such that $F_2(j) = \{0\}$.

It follows from definition that if E_1 is a cC_2 -space, then \check{E}_1 is an dC_2 -space. If E_2 is an dC_2 -space, then \check{E}_2 is a cC_2 -space.

Definition 8 We say that a C_2 -space (I_1, F_1, V_1) is a cfC_2 -space iff for any $i_1 \geq j_1 \in I_1$ the C_1 -space $F(i_1)/F(j_1)$ is a compact C_1 -space.

We say that a C_2 -space (I_2, F_2, V_2) is a dfC_2 -space iff for any $i_2 \geq j_2 \in I_2$ the C_1 -space $F(i_2)/F(j_2)$ is a discrete C_1 -space.

It follows from definition that if E_1 is a cfC_2 -space, then \check{E}_1 is a dfC_2 -space. If E_2 is a dfC_2 -space, then \check{E}_2 is a cfC_2 -space.

Definition 9 We say that a C_2 -space (I, F, V) is a complete C_2 -space if the following conditions are satisfied:

1. For any $i \geq j \in I$ the C_1 -space $E_{j,i} = F(i)/F(j)$ is a complete C_1 -space.
- 2.

$$V = \lim_{\substack{\longrightarrow \\ i \in I}} \lim_{\substack{\longleftarrow \\ j \leq i}} F(i)/F(j).$$

It follows from definitions that E is a complete C_2 -space iff $\check{\check{E}} = E$.

5.2 Virtual measures.

Let $E = (I, F, V)$ be a C_2 -space over the field \mathbb{F}_q .

For any $i, j \in I$ we define a 1-dimensional \mathbf{C} -vector space of virtual measures

$$\mu(F(i) \mid F(j)) \stackrel{\text{def}}{=} \varinjlim_{\substack{l \in I \\ l \leq i, l \leq j}} \text{Hom}_{\mathbf{C}}(\mu(F(i)/F(l)), \mu(F(j)/F(l))), \quad (91)$$

where to take the inductive limit we need the identities which follow from formula (27): for $l' \leq l \in I$

$$\begin{aligned} \mu(F(i)/F(l')) &= \mu(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(l)/F(l')) \\ \mu(F(j)/F(l')) &= \mu(F(j)/F(l)) \otimes_{\mathbf{C}} \mu(F(l)/F(l')). \end{aligned}$$

And

$$f \in \text{Hom}_{\mathbf{C}}(\mu(F(i)/F(l)), \mu(F(j)/F(l))) \mapsto f' \in \text{Hom}_{\mathbf{C}}(\mu(F(i)/F(l')), \mu(F(j)/F(l'))),$$

where $f'(a \otimes c) \stackrel{\text{def}}{=} f(a) \otimes c$, a is any from $\mu(F(i)/F(l))$, c is any from $\mu(F(l)/F(l'))$. This map is an isomorphism.

Proposition 22 *For any $i, j, l \in I$ there is a canonical isomorphism*

$$\gamma : \mu(F(i) \mid F(j)) \otimes_{\mathbf{C}} \mu(F(j) \mid F(l)) \longrightarrow \mu(F(i) \mid F(l))$$

such that the following diagram of associativity is commutative for any $i, j, l, n \in I$:

$$\begin{array}{ccc} \mu(F(i) \mid F(j)) \otimes_{\mathbf{C}} \mu(F(j) \mid F(l)) \otimes_{\mathbf{C}} \mu(F(l) \mid F(n)) & \longrightarrow & \mu(F(i) \mid F(l)) \otimes_{\mathbf{C}} \mu(F(l) \mid F(n)) \\ \downarrow & & \downarrow \\ \mu(F(i) \mid F(j)) \otimes_{\mathbf{C}} \mu(F(j) \mid F(n)) & \longrightarrow & \mu(F(i) \mid F(n)) \end{array}$$

Proof . We have a canonical map:

$$\begin{aligned} &\text{Hom}_{\mathbf{C}}(\mu(F(i)/F(l')), \mu(F(j)/F(l'))) \otimes_{\mathbf{C}} \text{Hom}_{\mathbf{C}}(\mu(F(j)/F(l')), \mu(F(l)/F(l'))) \\ &\quad \downarrow \\ &\text{Hom}_{\mathbf{C}}(\mu(F(i)/F(l')), \mu(F(l)/F(l'))) \end{aligned} \quad (92)$$

which satisfies the associativity diagram. And this map commutes with the inductive limit from the definition of $\mu(\cdot \mid \cdot)$. We obtain the map γ after the taking the inductive limit in (92).

Remark 10 We have the following canonical isomorphisms. For any $i, j \in I$

$$\mu(F(i) \mid F(i)) = \mathbf{C} \quad , \quad \mu(F(i) \mid F(j)) = \mu(F(j) \mid F(i))^*.$$

Let $i \leq j \in I$. Then

$$\mu(F(i) \mid F(j)) = \mu(F(j)/F(i)) \quad , \quad \mu(F(j) \mid F(i)) = \mu(F(j)/F(i))^*.$$

5.3 Basic spaces

Let $E = (I, F, V)$ be a C_2 -space over the field \mathbb{F}_q . Let $i \geq j \geq l \geq n \in I$. Then $F(i) \supset F(j) \supset F(l) \supset F(n)$, and from the definition of a C_2 -space we have

$$F(j)/F(n) \sim (F(i)/F(n)) \times_{F(i)/F(l)} (F(j)/F(l))$$

$$F(i)/F(l) \sim (F(i)/F(n)) \coprod_{F(j)/F(n)} (F(j)/F(l))$$

as C_1 -spaces.

For any $i \geq j \geq l \in I$ we have the following admissible triple of C_1 -spaces:

$$0 \rightarrow F(j)/F(l) \xrightarrow{\alpha_{lj}} F(i)/F(l) \xrightarrow{\beta_{li}} F(i)/F(j) \rightarrow 0.$$

We fix some $o \in I$. Then from section 4.7 we have that the following spaces over the field \mathbf{C} are well defined.

$$\begin{aligned} \mathcal{D}_{F(o)}(E) &\stackrel{\text{def}}{=} \lim_{\leftarrow i} \lim_{\leftarrow l} \mathcal{D}(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(l) \mid F(o)) = \\ &= \lim_{\leftarrow l} \lim_{\leftarrow i} \mathcal{D}(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(l) \mid F(o)) \end{aligned} \tag{93}$$

with respect to the maps $(\beta_{li})_* \otimes \mu(F(j) \mid F(o))$:

$$\mathcal{D}(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(l) \mid F(o)) \rightarrow \mathcal{D}(F(i)/F(j)) \otimes_{\mathbf{C}} \mu(F(j) \mid F(o)),$$

and the maps $\alpha_{li}^* \otimes \mu(F(l) \mid F(o))$:

$$\mathcal{D}(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(l) \mid F(o)) \rightarrow \mathcal{D}(F(j)/F(l)) \otimes_{\mathbf{C}} \mu(F(l) \mid F(o)).$$

$$\begin{aligned} \mathcal{D}'_{F(o)}(E) &\stackrel{\text{def}}{=} \lim_{\rightarrow i} \lim_{\rightarrow l} \mathcal{D}'(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(l)) = \\ &= \lim_{\rightarrow l} \lim_{\rightarrow i} \mathcal{D}'(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(l)) \end{aligned} \tag{94}$$

with respect to the maps $\beta_{li}^* \otimes \mu(F(o) \mid F(l))$:

$$\mathcal{D}'(F(i)/F(j)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(j)) \rightarrow \mathcal{D}'(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(l)),$$

and the maps $(\alpha_{li})_* \otimes \mu(F(o) \mid F(l))$:

$$\mathcal{D}'(F(j)/F(l)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(l)) \rightarrow \mathcal{D}'(F(i)/F(l)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(l)).$$

$$\mathcal{E}(E) \stackrel{\text{def}}{=} \lim_{\leftarrow i} \lim_{\leftarrow l} \mathcal{E}(F(i)/F(l)) \tag{95}$$

with respect to the maps α_{li}^* and β_{li}^* .

$$\tilde{\mathcal{E}}(E) \stackrel{\text{def}}{=} \lim_{\overrightarrow{l}} \lim_{\overleftarrow{i}} \tilde{\mathcal{E}}(F(i)/F(l)) \quad (96)$$

with respect to the maps β_{lji}^* and α_{lji}^* .

$$\mathcal{E}'(E) \stackrel{\text{def}}{=} \lim_{\overrightarrow{i}} \lim_{\overleftarrow{l}} \mathcal{E}'(F(i)/F(l)) \quad (97)$$

with respect to the maps $(\alpha_{lji})_*$ and $(\beta_{lji})_*$.

$$\tilde{\mathcal{E}}'(E) \stackrel{\text{def}}{=} \lim_{\overleftarrow{l}} \lim_{\overrightarrow{i}} \tilde{\mathcal{E}}'(F(i)/F(l)) \quad (98)$$

with respect to the maps $(\beta_{lji})_*$ and $(\alpha_{lji})_*$.

From definitions of these spaces we have the following canonical isomorphism for any $o_1 \in I$:

$$\mathcal{D}_{F(o)}(E) \otimes_{\mathbf{C}} \mu(F(o) \mid F(o_1)) \longrightarrow \mathcal{D}_{F(o_1)}(E)$$

such that the following diagram is commutative for any $o_2 \in I$:

$$\begin{array}{ccc} \mathcal{D}_{F(o)}(E) \otimes_{\mathbf{C}} \mu(F(o) \mid F(o_1)) \otimes_{\mathbf{C}} \mu(F(o_1) \mid F(o_2)) & \longrightarrow & \mathcal{D}_{F(o_1)}(E) \otimes_{\mathbf{C}} \mu(F(o_1) \mid F(o_2)) \\ \downarrow & & \downarrow \\ \mathcal{D}_{F(o)}(E) \otimes_{\mathbf{C}} \mu(F(o) \mid F(o_2)) & \longrightarrow & \mathcal{D}_{F(o_2)}(E) \end{array}$$

Dually, we have the following canonical isomorphism for any $o_1 \in I$:

$$\mu(F(o_1) \mid F(o)) \otimes_{\mathbf{C}} \mathcal{D}'_{F(o)}(E) \longrightarrow \mathcal{D}'_{F(o_1)}(E)$$

such that the following diagram is commutative for any $o_2 \in I$:

$$\begin{array}{ccc} \mu(F(o_2) \mid F(o_1)) \otimes_{\mathbf{C}} \mu(F(o_1) \mid F(o)) \otimes_{\mathbf{C}} \mathcal{D}'_{F(o)}(E) & \longrightarrow & \mu(F(o_2) \mid F(o_1)) \otimes_{\mathbf{C}} \mathcal{D}'_{F(o_1)}(E) \\ \downarrow & & \downarrow \\ \mu(F(o_2) \mid F(o)) \otimes_{\mathbf{C}} \mathcal{D}'_{F(o)}(E) & \longrightarrow & \mathcal{D}'_{F(o_2)}(E) \end{array}$$

Since for any $i \geq l \in I$ there are canonical embeddings

$$\begin{aligned} \tilde{\mathcal{E}}(F(i)/F(l)) &\hookrightarrow \mathcal{E}(F(i)/F(l)) \\ \mathcal{E}'(F(i)/F(l)) &\hookrightarrow \tilde{\mathcal{E}}'(F(i)/F(l)), \end{aligned}$$

there are the following canonical embeddings:

$$\begin{aligned} \tilde{\mathcal{E}}(E) &\hookrightarrow \mathcal{E}(E) \\ \mathcal{E}'(E) &\hookrightarrow \tilde{\mathcal{E}}'(E). \end{aligned}$$

Moreover, we have an embedding $\mathcal{E}(E) \longrightarrow \mathcal{F}(V)$ such that $\tilde{\mathcal{E}}(E)$ and $\mathcal{E}(E)$ are \mathbf{C} -subalgebras of \mathbf{C} -algebra $\mathcal{F}(V)$.

Proposition 23 *The following statements are satisfied.*

1. The \mathbf{C} -vector space $\mathcal{D}_{F(o)}(E)$ is a module over the \mathbf{C} -algebra $\mathcal{E}(E)$.
2. The \mathbf{C} -vector space $\mathcal{D}'_{F(o)}(E)$ is a module over the \mathbf{C} -algebra $\mathcal{E}(E)$.

Proof . We construct the map:

$$\mathcal{E}(E) \otimes_{\mathbf{C}} \mathcal{D}_{F(o)}(E) \longrightarrow \mathcal{D}_{F(o)}(E). \quad (99)$$

For any $i_1 \geq j_1 \in I$ we have the following multiplication map:

$$\mathcal{E}(F(i_1)/F(j_1)) \otimes_{\mathbf{C}} \mathcal{D}(F(i_1)/F(j_1)) \otimes_{\mathbf{C}} \mu(F(j_1)|F(o)) \xrightarrow{\lambda} \mathcal{D}(F(i_1)/F(j_1)) \otimes_{\mathbf{C}} \mu(F(j_1)|F(o)). \quad (100)$$

Now from projection formula (40) we have the following formula for any $i \geq j \geq l \in I$, for any $g \in \mathcal{E}(F(i)/F(j))$, $f \in \mathcal{D}(F(i)/F(l))$, $\mu \in \mu(F(l) | F(o))$:

$$((\beta_{lji})_* \otimes \mu(F(j)/F(o))) (\lambda(\beta_{lji}^*(g) \otimes f \otimes \mu)) = \lambda(g \otimes ((\beta_{lji})_* \otimes \mu(F(j)/F(o))) (f \otimes \mu)). \quad (101)$$

Using formulas (95) and (93) we construct map (99) from maps (100). Due to formulas (101) it is well defined. We proved the first statement of the proposition.

For any $i_1 \geq j_1 \in I$, any $f \in \mathcal{E}(F(i_1)/F(j_1))$ we defined in remark 5 the action of f on $\mathcal{D}'(F(i_1)/F(j_1))$ as conjugate action to the action of f on $\mathcal{D}(F(i_1)/F(j_1))$. Therefore applying the conjugate formulas to the formulas above, from formulas (95) and (94) we construct the conjugate action of $\mathcal{E}(E)$ on $\mathcal{D}'_{F(o)}(E)$.

Remark 11 Let $\{B_l\}$, $l \in J$ be a projective system of vector spaces over a field k with the surjective transition maps $\phi_{l_1, l_2} : B_{l_1} \rightarrow B_{l_2}$ for any $l_1 \geq l_2 \in J$. Let J^0 be a partially ordered set with the same set as J but with inverse order then J . Let $\{A_l\}$, $l \in J^0$ be an inductive system of k -vector spaces with the injective transition maps $\psi_{l_2, l_1} : A_{l_2} \rightarrow A_{l_1}$. Suppose for any $l \in J$ we have a nondegenerate k -linear pairing

$$\langle \cdot, \cdot \rangle_l : B_l \times A_l \longrightarrow k$$

such that for any $l_1 \geq l_2 \in I$, $x \in B_{l_1}$, $y \in A_{l_2}$ we have

$$\langle \phi_{l_1, l_2}(x), y \rangle_{l_2} = \langle x, \psi_{l_2, l_1}(y) \rangle_{l_1}.$$

Then we have canonically nondegenerate k -linear pairing between k -vector spaces $\varprojlim_{l \in J} B_l$ and $\varinjlim_{l \in J^0} A_l$ which is induced by pairings $\langle \cdot, \cdot \rangle_l$, $l \in J$.

For a C_2 -space $E = (I, F, V)$ over the field \mathbb{F}_q , for any $i \geq j \in I$ there are nondegenerate \mathbf{C} -linear pairings between $\mathcal{D}(F(i)/F(j))$ and $\mathcal{D}'(F(i)/F(j))$, between $\mathcal{E}(F(i)/F(j))$ and $\mathcal{E}'(F(i)/F(j))$, between $\mathcal{E}(F(i)/F(j))$ and $\mathcal{E}'(F(i)/F(j))$ (see remark 4).

Therefore applying remark 11 twice to formulas (93)-(98) we obtain that there are the following nondegenerate pairings:

$$\langle \cdot, \cdot \rangle_{\mathcal{D}_{F(o)}(E)} : \mathcal{D}_{F(o)}(E) \times \mathcal{D}'_{F(o)}(E) \longrightarrow \mathbf{C}$$

$$\begin{aligned} \langle \cdot, \cdot \rangle_{\mathcal{E}(E)} &: \mathcal{E}(E) \times \mathcal{E}'(E) \rightarrow \mathbf{C} \\ \langle \cdot, \cdot \rangle_{\tilde{\mathcal{E}}(E)} &: \tilde{\mathcal{E}}'(E) \times \tilde{\mathcal{E}}(E) \rightarrow \mathbf{C}. \end{aligned}$$

Now from the construction of multiplications from proposition 23 we have for any $g \in \mathcal{E}(E)$, $f \in \mathcal{D}_{F(o)}(E)$, $G \in \mathcal{D}'_{F(o)}(E)$:

$$\langle f, g \cdot G \rangle_{\mathcal{D}_{F(o)}(E)} = \langle g \cdot f, G \rangle_{\mathcal{D}_{F(o)}(E)}. \quad (102)$$

Remark 12 Let $E = (I, F, V)$ be a cfC_2 -space over the field \mathbb{F}_q . Then for any $i \geq j \in I$ we have a canonical element

$$1_{ij} \in \mu(F(i)/F(j)) \quad : \quad 1_{ij}(\delta_{(F(i)/F(j))}) \stackrel{\text{def}}{=} 1.$$

Therefore for any $k, l \in I$ we have a canonical element $1_{kl} \in \mu(F(k) \mid F(l))$ such that for any $k, l, n \in I$ according to proposition 22:

$$\gamma(1_{kl} \otimes 1_{ln}) = 1_{kn}.$$

Taking into account these elements 1_{lo} and 1_{ol} for $l, o \in I$ we can do not write 1-dimensional \mathbf{C} -spaces $\mu(F(l) \mid F(o))$ and $\mu(F(o) \mid F(l))$ in formulas (93) and (94) which define \mathbf{C} -spaces $\mathcal{D}_{F(o)}(E)$ and $\mathcal{D}'_{F(o)}(E)$. These \mathbf{C} -spaces *do not depend* on the choice of $o \in I$.

Similarly, Let $E = (I, F, V)$ be a dfC_2 -space over the field \mathbb{F}_q . Then for any $i \geq j \in I$ we have a canonical element

$$\delta_{ij} \in \mu(F(i)/F(j)) \quad : \quad \delta_{ij}(\{0\}) \stackrel{\text{def}}{=} 1.$$

Therefore for any $k, l \in I$ we have a canonical element $\delta_{kl} \in \mu(F(k) \mid F(l))$ such that for any $k, l, n \in I$ according to proposition 22:

$$\gamma(\delta_{kl} \otimes \delta_{ln}) = \delta_{kn}.$$

Taking into account these elements δ_{lo} and δ_{ol} for $l, o \in I$ we can do not write 1-dimensional \mathbf{C} -spaces $\mu(F(l) \mid F(o))$ and $\mu(F(o) \mid F(l))$ in formulas (93) and (94) which define \mathbf{C} -spaces $\mathcal{D}_{F(o)}(E)$ and $\mathcal{D}'_{F(o)}(E)$. These \mathbf{C} -spaces *do not depend* on the choice of $o \in I$.

Remark 13 From a C_2 -space $E = (I, F, V)$ over the field \mathbb{F}_q , we constructed 6 \mathbf{C} -spaces: $\mathcal{D}_{F(o)}(E)$, $\mathcal{D}'_{F(o)}(E)$ (for any $o \in I$), $\mathcal{E}(E)$, $\tilde{\mathcal{E}}(E)$, $\mathcal{E}'(E)$, $\tilde{\mathcal{E}}'(E)$ by formulas (93)-(98).

We can construct 10 \mathbf{C} -spaces, where additional \mathbf{C} -spaces appear if for any $i \geq l \in I$ we put $\tilde{\mathcal{E}}(F(i)/F(l))$ instead $\mathcal{E}(F(i)/F(l))$ in formula (95), $\mathcal{E}(F(i)/F(l))$ instead $\tilde{\mathcal{E}}(F(i)/F(l))$ in formula (96), $\mathcal{E}'(F(i)/F(l))$ instead $\tilde{\mathcal{E}}'(F(i)/F(l))$ in formula (98), $\tilde{\mathcal{E}}'(F(i)/F(l))$ instead $\mathcal{E}'(F(i)/F(l))$ in formula (97).

In the sequel we restrict ourselves to consider 6 \mathbf{C} -spaces which are given by formulas (93)-(98).

5.4 Fourier transform

Let $E = (I, F, V)$ be a complete C_2 -space over the field \mathbb{F}_q . Then we have $\check{\check{E}} = E$. We fix some $o \in I$.

5.4.1

Let $i \geq j \in I$ be any. By section 4.4 there are the following \mathbf{C} -linear maps:

$$f \in \mathcal{E}(F(i)/F(j)) \longmapsto \check{f} \in \mathcal{E}(F(i)/F(j))$$

$$G \in \tilde{\mathcal{E}}'(F(i)/F(j)) \longmapsto \check{G} \in \tilde{\mathcal{E}}'(F(i)/F(j))$$

$$\begin{aligned} f \otimes \mu &\in \mathcal{D}(F(i)/F(j)) \otimes_{\mathbf{C}} \mu(F(j) \mid F(o)) \mapsto \check{f} \otimes \mu \in \mathcal{D}(F(i)/F(j)) \otimes_{\mathbf{C}} \mu(F(j) \mid F(o)) \\ G \otimes \mu' &\in \mathcal{D}'(F(i)/F(j)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(j)) \mapsto \check{G} \otimes \mu' \in \mathcal{D}'(F(i)/F(j)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(j)). \end{aligned}$$

They map the \mathbf{C} -subspaces $\tilde{\mathcal{E}}(F(i)/F(j)) \subset \mathcal{E}(F(i)/F(j))$ and $\mathcal{E}'(F(i)/F(j)) \subset \tilde{\mathcal{E}}'(F(i)/F(j))$ to itself correspondingly.

These maps commute with direct and inverse images when we change $i, j \in I$ to $i', j' \in I$, $i' \geq i$, $j' \leq j$. Therefore using formulas (93)-(98) we have that the following \mathbf{C} -linear maps are well defined:

$$f \in \mathcal{D}_{F(o)}(E) \longmapsto \check{f} \in \mathcal{D}_{F(o)}(E)$$

$$G \in \mathcal{D}'_{F(o)}(E) \longmapsto \check{G} \in \mathcal{D}'_{F(o)}(E)$$

$$f \in \mathcal{E}(E) \longmapsto \check{f} \in \mathcal{E}(E)$$

$$G \in \tilde{\mathcal{E}}'(E) \longmapsto \check{G} \in \tilde{\mathcal{E}}'(E).$$

They map the \mathbf{C} subspaces $\tilde{\mathcal{E}}(E) \subset \mathcal{E}(E)$ and $\mathcal{E}'(E) \subset \tilde{\mathcal{E}}'(E)$ to itself correspondingly.

The square of any from these maps is the identity map. From the corresponding formulas in 1-dimensional case we have the following formulas:

$$\begin{aligned} \langle \check{f}, G \rangle_{\mathcal{D}_{F(o)}(E)} &= \langle f, \check{G} \rangle_{\mathcal{D}_{F(o)}(E)} \quad \text{for any } f \in \mathcal{D}_{F(o)}(E), \quad G \in \mathcal{D}'_{F(o)}(E); \\ \langle \check{f}, G \rangle_{\mathcal{E}(E)} &= \langle f, \check{G} \rangle_{\mathcal{E}(E)} \quad \text{for any } f \in \mathcal{E}(E), \quad G \in \mathcal{E}'(E); \\ \langle \check{G}, f \rangle_{\tilde{\mathcal{E}}(E)} &= \langle G, \check{f} \rangle_{\tilde{\mathcal{E}}(E)} \quad \text{for any } G \in \tilde{\mathcal{E}}'(E), \quad f \in \tilde{\mathcal{E}}(E). \end{aligned}$$

5.4.2

According to section 5.1 we have the C_2 -space $\check{E} = (I^0, F^0, \check{V})$. For any $i \geq j \in I$, for the C_1 -space $E_{j,i} = F(i)/F(j)$ we have the C_1 -space $\check{E}_{j,i} = F^0(j)/F^0(i)$.

For any $l, n \in I$ we have

$$\mu(F(l) \mid F(n)) = \mu(F^0(l) \mid F^0(n)). \quad (103)$$

Indeed, let $k \in I$ such that $k \leq l$, $k \leq n$. Then according to proposition 22, remark 10 and formula (23) we have

$$\begin{aligned} \mu(F(l) \mid F(n)) &= \mu(F(l) \mid F(k)) \otimes_{\mathbf{C}} \mu(F(k) \mid F(n)) = \\ &= \mu(F(l)/F(k))^* \otimes_{\mathbf{C}} \mu(F(n)/F(k)) = \mu(F^0(k)/F^0(l)) \otimes_{\mathbf{C}} \mu(F^0(k)/F^0(n))^* = \\ &= \mu(F^0(l) \mid F^0(k)) \otimes_{\mathbf{C}} \mu(F^0(k) \mid F^0(n)) = \mu(F^0(l) \mid F^0(n)). \end{aligned}$$

Therefore we have the following maps for any $i \geq j \in I$:

$$\begin{aligned} \mathbf{F} \otimes \mu(F(i) \mid F(o)) & : \\ \mathcal{D}(F(i)/F(j)) \otimes_{\mathbf{C}} \mu(F(j) \mid F(o)) & \longrightarrow \mathcal{D}(F^0(j)/F^0(i)) \otimes_{\mathbf{C}} \mu(F^0(i) \mid F^0(o)) \end{aligned} \quad (104)$$

$$\begin{aligned} \mathbf{F} \otimes \mu(F(o) \mid F(i)) & : \\ \mathcal{D}'(F(i)/F(j)) \otimes_{\mathbf{C}} \mu(F(o) \mid F(j)) & \longrightarrow \mathcal{D}'(F^0(j)/F^0(i)) \otimes_{\mathbf{C}} \mu(F^0(o) \mid F^0(i)) \end{aligned} \quad (105)$$

$$\mathbf{F} : \mathcal{E}(F(i)/F(j)) \longrightarrow \tilde{\mathcal{E}}'(F^0(j)/F^0(i)) \quad (106)$$

$$\mathbf{F} : \tilde{\mathcal{E}}'(F(i)/F(j)) \longrightarrow \mathcal{E}(F^0(j)/F^0(i)) \quad (107)$$

$$\mathbf{F} : \tilde{\mathcal{E}}(F(i)/F(j)) \longrightarrow \mathcal{E}'(F^0(j)/F^0(i)) \quad (108)$$

$$\mathbf{F} : \mathcal{E}'(F(i)/F(j)) \longrightarrow \tilde{\mathcal{E}}(F^0(j)/F^0(i)). \quad (109)$$

We remark that the map \mathbf{F} from formula (106) being restricted to the \mathbf{C} -subspace $\tilde{\mathcal{E}}(F(i)/F(j)) \subset \mathcal{E}(F(i)/F(j))$ coincides with the map \mathbf{F} from formula (108). The map \mathbf{F} from formula (107) being restricted to the \mathbf{C} -subspace $\mathcal{E}'(F(i)/F(j)) \subset \tilde{\mathcal{E}}'(F(i)/F(j))$ coincides with the map \mathbf{F} from formula (109).

Now we use diagrams (72)-(75), (77)-(80), which connect Fourier transform with direct and inverse images. We use it when we change $i, j \in I$ to $i', j' \in I$, $i' \geq i$, $j' \leq j$, and take the limits according to formulas (93)-(98). We obtain that the following \mathbf{C} -linear maps (*two-dimensional Fourier transforms*) are well defined:

$$\mathbf{F} : \mathcal{D}_{F(o)}(E) \longrightarrow \mathcal{D}_{F^0(o)}(\check{E}) \quad (110)$$

$$\mathbf{F} : \mathcal{D}'_{F(o)}(E) \longrightarrow \mathcal{D}'_{F^0(o)}(\check{E}) \quad (111)$$

$$\mathbf{F} : \mathcal{E}(E) \longrightarrow \tilde{\mathcal{E}}'(\check{E}) \quad (112)$$

$$\mathbf{F} : \tilde{\mathcal{E}}'(E) \longrightarrow \mathcal{E}(\check{E}) \quad (113)$$

$$\mathbf{F} : \tilde{\mathcal{E}}(E) \longrightarrow \mathcal{E}'(\check{E}) \quad (114)$$

$$\mathbf{F} : \mathcal{E}'(E) \longrightarrow \tilde{\mathcal{E}}(\check{E}). \quad (115)$$

We remark that from the construction and the corresponding one-dimensional property it follows that the map \mathbf{F} from formula (112) being restricted to the \mathbf{C} -subspace $\tilde{\mathcal{E}}(E) \subset \mathcal{E}(E)$ coincides with the map \mathbf{F} from formula (114). The map \mathbf{F} from formula (113) being restricted to the \mathbf{C} -subspace $\mathcal{E}'(E) \subset \tilde{\mathcal{E}}'(E)$ coincides with the map \mathbf{F} from formula (115).

Proposition 24 *The following statements are satisfied.*

1. *The maps \mathbf{F} are isomorphisms of \mathbf{C} -vector spaces from formulas (110)-(115).*
2. *For any $f \in \mathcal{D}_{F(o)}(E)$ and $G \in \mathcal{D}'_{F(o)}(E)$*

$$\mathbf{F} \circ \mathbf{F}(f) = \check{f}, \quad \mathbf{F} \circ \mathbf{F}(G) = \check{G},$$

for any $f \in \mathcal{E}(E)$ and $G \in \tilde{\mathcal{E}}'(E)$

$$\mathbf{F} \circ \mathbf{F}(f) = \check{f}, \quad \mathbf{F} \circ \mathbf{F}(G) = \check{G};$$

3. For any $f \in \mathcal{D}_{F(o)}(E)$ and $G \in \mathcal{D}'_{F(o)}(E)$

$$\langle \mathbf{F}(f), G \rangle_{\mathcal{D}_{F(o)}(E)} = \langle f, \mathbf{F}(G) \rangle_{\mathcal{D}_{F(o)}(E)} ,$$

for any $f \in \mathcal{E}(E)$ and $g \in \tilde{\mathcal{E}}(E)$

$$\langle \mathbf{F}(f), g \rangle_{\tilde{\mathcal{E}}(E)} = \langle f, \mathbf{F}(g) \rangle_{\mathcal{E}(E)} ,$$

for any $H \in \tilde{\mathcal{E}}'(E)$, $G \in \mathcal{E}'(E)$

$$\langle \mathbf{F}(H), G \rangle_{\mathcal{E}(E)} = \langle H, \mathbf{F}(G) \rangle_{\tilde{\mathcal{E}}(E)} .$$

Proof follows from the construction of two-dimensional Fourier transform given in this section and the properties of 1-dimensional Fourier transform (see section 4.4) applied to functions and distributions on C_1 -spaces $F(i)/F(j)$ for any $i \geq j \in I$.

5.5 Central extension and its representations

In this section we construct a central extension of group of automorphisms of an object of the category C_2 and consider representations of this group on the spaces of functions and distributions of the given object.

5.5.1

Let a C_2 -space $E_1 = (I_1, F_1, V)$ dominates a C_2 -space $E_2 = (I_2, F_2, V)$ (over the field \mathbb{F}_q). We fix some $o \in I_2$. Then we obtain the following canonical isomorphisms:

$$\begin{aligned} \mathcal{D}_{F(o)}(E_1) &= \mathcal{D}_{F(o)}(E_2), & \mathcal{D}'_{F(o)}(E_1) &= \mathcal{D}'_{F(o)}(E_2), \\ \mathcal{E}(E_1) &= \mathcal{E}(E_2), & \mathcal{E}'(E_1) &= \mathcal{E}'(E_2), \\ \tilde{\mathcal{E}}(E_1) &= \tilde{\mathcal{E}}(E_2), & \tilde{\mathcal{E}}'(E_1) &= \tilde{\mathcal{E}}'(E_2). \end{aligned}$$

Indeed, from formulas (93)-(98) we obtain that inductive and projective limits used in the definitions of these spaces will be the same on sets of indices depending on I_1 or I_2 . Moreover, by the same reasons, the two-dimensional Fourier transform coincide for the spaces depending on E_1 or E_2 . (Compare it with section 4.9.)

5.5.2

Let $E = (I, F, V)$ be a C_2 -space over a field k . We construct the maximal C_2 -space \tilde{E} over the field k , which dominates the C_2 -space E . By $G(E)$ we denote the set of all k -subspaces $W \subset V$ such that there are some $i \geq j \in I$ with the properties:

- $F(i) \supset W \supset F(j)$,
- for any $x \in F(i)/F(j)$, $x \notin W/F(j)$ there is $l \in I_{ij}$ such that $(x + F_{ij}(l)) \cap (W/F(j)) = \emptyset$ (this intersection is inside $F(i)/F(j)$), where $(I_{ij}, F_{ij}, F(i)/F(j))$ is the structure of C_1 -space on the k -space $F(i)/F(j)$.

Then $G(E)$ is a partially ordered set, which is ordered by inclusions of subspaces. We remark that for any $k, l \in G(E)$ there are $p_1 \geq p_2 \in G(E)$ such that $p_1 \geq k \geq p_2$, $p_1 \geq l \geq p_2$. Therefore V is a filtered k -vector space with filtration given by $G(E)$ (see section 4.1).

For any $W_1, W_2 \in G(E)$ such that $W_1 \supset W_2$ we introduce the structure of C_1 -space on the k -vector space W_1/W_2 in the following way. There are $i \geq j \in I$ such that

$$F(i) \supset W_1 \supset W_2 \supset F(j).$$

The k -space $F(i)/F(j)$ is a C_1 -space, therefore it is filtered. We restrict the filtration from the k -space $F(i)/F(j)$ to the k -space $W_1/F(j)$. After that we take the factor filtration on the k -space W_1/W_2 . By $G_{i,j}(W_1/W_2)$ we denote the set of k -subspaces of W_1/W_2 which are given by this filtration. Now by $Gr(W_1/W_2)$ we denote the set of all k -subspaces $U \subset W_1/W_2$ such that there are some $P, Q \in G_{i,j}(W_1/W_2)$ with the property

$$P \supset U \supset Q.$$

Then $Gr(W_1/W_2)$ is a partially ordered set, which is ordered by inclusions of subspaces. The filtration on W_1/W_2 which is given by elements of $Gr(W_1/W_2)$ defines a well-defined C_1 -space on W_1/W_2 .

Thus we constructed the well-defined C_2 -space \tilde{E} , which is filtered by the set $G(E)$. We denote this C_2 -space by $(\tilde{I}, \tilde{F}, V)$.

5.5.3

Let $E = (I, F, V)$ be a C_2 -space over the field \mathbb{F}_q . Then for any $i_1, i_2 \in \tilde{I}$ we have the virtual measures space $\mu(\tilde{F}(i_1) | \tilde{F}(i_2))$. We have $\text{Aut}_{C_2}(E) = \text{Aut}_{C_2}(\tilde{E})$. For any $g \in \text{Aut}_{C_2}(E)$, for any $i \in I$ we have from the definition of the group $\text{Aut}_{C_2}(E)$ that $gF(i) = \tilde{F}(p)$ for some $p \in \tilde{I}$. Therefore for any $g, h \in \text{Aut}_{C_2}(E)$, for any $i_1, i_2 \in I$ we have the well-defined 1-dimensional \mathbf{C} -vector space $\mu(gF(i_1) | hF(i_2))$.

For any $g \in \text{Aut}_{C_2}(E)$, for any $p \geq q \in \tilde{I}$ we have the following \mathbf{C} -isomorphism

$$n_g : \mu(\tilde{F}(p)/\tilde{F}(q)) \longrightarrow \mu(g\tilde{F}(p)/g\tilde{F}(q)),$$

where for any $U \in Gr(g\tilde{F}(p)/g\tilde{F}(q))$, for any $\mu \in \mu(\tilde{F}(p)/\tilde{F}(q))$:

$$n_g(\mu)(\delta_U) \stackrel{\text{def}}{=} \mu(\delta_{g^{-1}U}).$$

For any $g \in \text{Aut}_{C_2}(E)$, for any $p, q, s \in \tilde{I}$, $s \leq p$, $s \leq q$ we have the following \mathbf{C} -isomorphism m_g :

$$\text{Hom}_{\mathbf{C}}(\mu(\tilde{F}(p)/\tilde{F}(s)), \mu(\tilde{F}(q)/\tilde{F}(s))) \longrightarrow \text{Hom}_{\mathbf{C}}(\mu(g\tilde{F}(p)/g\tilde{F}(s)), \mu(g\tilde{F}(q)/g\tilde{F}(s))),$$

where for any $f \in \text{Hom}_{\mathbf{C}}(\mu(\tilde{F}(p)/\tilde{F}(s)), \mu(\tilde{F}(q)/\tilde{F}(s)))$:

$$m_g(f) \stackrel{\text{def}}{=} n_g \circ f \circ n_{g^{-1}}.$$

No we apply isomorphisms m_g to the inductive limit in formula (91), then we obtain the following \mathbf{C} -linear isomorphism for any $p, q \in \tilde{I}$:

$$l_g : \mu(\tilde{F}(p) \mid \tilde{F}(q)) \longrightarrow \mu(g\tilde{F}(p) \mid g\tilde{F}(q)).$$

We have for any $g_1, g_2 \in G$ that $l_{g_1 g_2} = l_{g_1} l_{g_2}$. Besides, $\gamma(l_g(a) \otimes l_g(b)) = l_g \gamma(a \otimes b)$ (see proposition 22) for any $a \in \mu(\tilde{F}(p) \mid \tilde{F}(q))$, $b \in \mu(\tilde{F}(q) \mid \tilde{F}(s))$, for any $p, q, s \in \tilde{I}$.

Let $p, q \in \tilde{I}$. Then for any $\mu \in \mu(\tilde{F}(p) \mid \tilde{F}(q))$, $\mu \neq 0$ we define canonically $\mu^{-1} \in \mu(\tilde{F}(q) \mid \tilde{F}(p))$ such that $\mu \otimes \mu^{-1} = 1$ with respect to the following canonical isomorphism:

$$\mu(\tilde{F}(p) \mid \tilde{F}(q)) \otimes_{\mathbf{C}} \mu(\tilde{F}(q) \mid \tilde{F}(p)) = \mathbf{C}.$$

Let $E = (I, F, V)$ be a C_2 -space over the field \mathbb{F}_q . We fix some $o \in I$. Then there is the following central extension of groups:

$$1 \longrightarrow \mathbf{C}^* \longrightarrow \widehat{\text{Aut}_{C_2}(E)}_{F(o)} \xrightarrow{\Lambda} \text{Aut}_{C_2}(E) \longrightarrow 1, \quad (116)$$

where

$$\widehat{\text{Aut}_{C_2}(E)}_{F(o)} \stackrel{\text{def}}{=} \{(g, \mu) : g \in \text{Aut}_{C_2}(E), \mu \in \mu(F(o) \mid gF(o)), \mu \neq 0\}.$$

Here $\Lambda((g, \mu)) = g$. The operations $(g_1, \mu_1) \cdot (g_2, \mu_2) = (g_1 g_2, \gamma(\mu_1 \otimes l_{g_1}(\mu_2)))$ and $(g, \mu)^{-1} = (g^{-1}, l_{g^{-1}}(\mu^{-1}))$ define the structure of a group on the set $\widehat{\text{Aut}_{C_2}(E)}_{F(o)}$. (The unit element of this group is $(e, 1)$, where e is the unit element of the group $\text{Aut}_{C_2}(E)$).

Remark 14 For any $o_1 \in I$ there is a canonical isomorphism

$$\alpha_{o, o_1} : \widehat{\text{Aut}_{C_2}(E)}_{F(o)} \longrightarrow \widehat{\text{Aut}_{C_2}(E)}_{F(o_1)}.$$

Indeed, we fix any $\nu \in \mu(F(o_1) \mid F(o))$, $\nu \neq 0$. Then

$$\alpha_{o, o_1}((g, \mu)) \stackrel{\text{def}}{=} (g, \gamma(\gamma(\nu \otimes \mu) \otimes l_g(\nu^{-1}))).$$

The map α_{o, o_1} does not depend on the choice of $\nu \in \mu(F(o_1) \mid F(o))$, $\nu \neq 0$.

Remark 15 For any C_2 -space $E = (I, F, V)$ over a field k , any $o \in I$ it is possible to construct the central extension:

$$0 \longrightarrow \mathbb{Z} \longrightarrow \widetilde{\text{Aut}_{C_2}(E)}_{F(o)} \longrightarrow \text{Aut}_{C_2}(E) \longrightarrow 1. \quad (117)$$

If the field $k = \mathbb{F}_q$, then under the map

$$a \in \mathbb{Z} \longmapsto q^a \in \mathbf{C}^*,$$

which we apply to the kernel of central extension (117), central extension (117) transfers to central extension (116).

In the case if a C_2 -space E is constructed from the two-dimensional local field $k((t_1))((t_2))$, then central extension (117) was constructed in [11].

5.5.4

Let $E = (I, F, V)$ be a C_2 -space over the finite field \mathbb{F}_q . In section 5.5.2 we constructed the maximal C_2 -space $\tilde{E} = (\tilde{I}, \tilde{F}, V)$ which dominates the C_2 -space E . We recall that $G(E)$ is the set of all k -subspaces $W \subset V$ such that $W = \tilde{F}(p)$ for some $p \in \tilde{I}$. Then any $g \in \text{Aut}_{C_2}(E)$ induces an isomorphism of the set $G(E)$.

For any $g \in \text{Aut}_{C_2}(E)$, for any $p \geq q \in \tilde{I}$ we have the following maps:

$$r_g : \mathcal{D}(\tilde{F}(p)/\tilde{F}(q)) \longrightarrow \mathcal{D}(g\tilde{F}(p)/g\tilde{F}(q)) \quad (118)$$

$$r_g : \tilde{\mathcal{E}}(\tilde{F}(p)/\tilde{F}(q)) \longrightarrow \tilde{\mathcal{E}}(g\tilde{F}(p)/g\tilde{F}(q)) \quad (119)$$

$$r_g : \mathcal{E}(\tilde{F}(p)/\tilde{F}(q)) \longrightarrow \mathcal{E}(g\tilde{F}(p)/g\tilde{F}(q)), \quad (120)$$

where $r_g(f)(v) \stackrel{\text{def}}{=} f(g^{-1}v)$ for any $v \in g\tilde{F}(p)/g\tilde{F}(q)$ and any f from $\mathcal{D}(\tilde{F}(p)/\tilde{F}(q))$, or any f from $\tilde{\mathcal{E}}(g\tilde{F}(p)/g\tilde{F}(q))$, or any f from $\mathcal{E}(\tilde{F}(p)/\tilde{F}(q))$.

Applying formulas (119)-(120) to formulas (96) and (95) we obtain the maps for any $g \in \text{Aut}_{C_2}(E)$

$$r_g : \tilde{\mathcal{E}}(\tilde{E}) \longrightarrow \tilde{\mathcal{E}}(\tilde{E})$$

$$r_g : \mathcal{E}(\tilde{E}) \longrightarrow \mathcal{E}(\tilde{E})$$

such that $r_g r_h = r_{gh}$ for any $g, h \in \text{Aut}_{C_2}(E)$.

Therefore using section 5.5.1 we obtain the representations of the group $\text{Aut}_{C_2}(E)$ by maps r_g on the \mathbf{C} -spaces $\tilde{\mathcal{E}}(E)$ and $\mathcal{E}(E)$ such that the first one is a subrepresentation of the second one.

We fix some $o \in I$. Using formula (118) we construct the map $R_{\tilde{g}}$ for any $\tilde{g} = (g, \mu) \in \widehat{\text{Aut}_{C_2}(E)}_{F(o)}$ for any $p \geq q \in \tilde{I}$:

$$R_{\tilde{g}} : \mathcal{D}(\tilde{F}(p)/\tilde{F}(q)) \otimes_{\mathbf{C}} \mu(\tilde{F}(q) \mid F(o)) \longrightarrow \mathcal{D}(g\tilde{F}(p)/g\tilde{F}(q)) \otimes_{\mathbf{C}} \mu(g\tilde{F}(q) \mid F(o)) \quad (121)$$

as composition of the map $r_g \otimes l_g$ with multiplication by $\mu^{-1} \in \mu(gF(o) \mid F(o))$.

Applying this formula to formula (93) we obtain the map for any $\tilde{g} \in \widehat{\text{Aut}_{C_2}(E)}_{F(o)}$:

$$R_{\tilde{g}} : \mathcal{D}_{F(o)}(\tilde{E}) \longrightarrow \mathcal{D}_{F(o)}(\tilde{E})$$

such that $R_{\tilde{g}} R_{\tilde{h}} = R_{\tilde{g}\tilde{h}}$.

Therefore using section 5.5.1 we obtain the representation of the group $\widehat{\text{Aut}_{C_2}(E)}_{F(o)}$ by maps $R_{\tilde{g}}$ on the \mathbf{C} -space $\mathcal{D}_{F(o)}(E)$.

From constructions we have the following formula:

$$R_{\tilde{g}}(f \cdot H) = r_{\Lambda(\tilde{g})}(f) \cdot R_{\tilde{g}}(H) \quad (122)$$

for any $\tilde{g} \in \widehat{\text{Aut}_{C_2}(E)}_{F(o)}$, $f \in \mathcal{E}(E)$, $H \in \mathcal{D}(E)$.

Dually to formulas (119)-(120), for any $g \in \text{Aut}_{C_2}(E)$, for any $p \geq q \in \tilde{I}$ we have the following maps:

$$r'_g : \mathcal{E}'(\tilde{F}(p)/\tilde{F}(q)) \longrightarrow \mathcal{E}'(g\tilde{F}(p)/g\tilde{F}(q)), \quad (123)$$

$$r'_g : \tilde{\mathcal{E}}'(\tilde{F}(p)/\tilde{F}(q)) \longrightarrow \tilde{\mathcal{E}}'(g\tilde{F}(p)/g\tilde{F}(q)). \quad (124)$$

Applying formulas (123)-(124) to formulas (97)-(98), and using section 5.5.1, we obtain the representations of the group $\text{Aut}_{C_2}(E)$ on the \mathbf{C} -spaces $\tilde{\mathcal{E}}'(E)$ and $\mathcal{E}'(E)$ by maps r'_g such that the first representation is a subrepresentation of the second one. By construction, we have for any $f_1 \in \mathcal{E}(E)$, any $H_1 \in \mathcal{E}'(E)$, and any $g \in \text{Aut}_{C_2}(E)$

$$\langle r_g f_1, r'_g H_1 \rangle_{\mathcal{E}(E)} = \langle f_1, H_1 \rangle_{\mathcal{E}(E)}.$$

We have for any $f_2 \in \tilde{\mathcal{E}}(E)$, any $H_2 \in \tilde{\mathcal{E}}'(E)$, and any $g \in \text{Aut}_{C_2}(E)$

$$\langle r'_g H_2, r_g f_2 \rangle_{\tilde{\mathcal{E}}(E)} = \langle H_2, f_2 \rangle_{\tilde{\mathcal{E}}(E)}.$$

Let $o \in I$. Dually to formula (121) we construct the map R'_g for any $\tilde{g} = (g, \mu) \in \widehat{\text{Aut}_{C_2}(E)}_{F(o)}$ for any $p \geq q \in \tilde{I}$:

$$R'_g : \mathcal{D}'(\tilde{F}(p)/\tilde{F}(q)) \otimes_{\mathbf{C}} \mu(F(o) \mid \tilde{F}(q)) \longrightarrow \mathcal{D}'(g\tilde{F}(p)/g\tilde{F}(q)) \otimes_{\mathbf{C}} \mu(F(o) \mid g\tilde{F}(q)). \quad (125)$$

Applying formula (125) to formula (94), and using section 5.5.1, we obtain the representation of the group $\widehat{\text{Aut}_{C_2}(E)}_{F(o)}$ on the \mathbf{C} -space $\mathcal{D}'_{F(o)}(E)$ by maps R'_g . By construction, we have the following formula for any $H \in \mathcal{D}'_{F(o)}(E)$, any $f \in \mathcal{D}_{F(o)}(E)$, any $\tilde{g} \in \widehat{\text{Aut}_{C_2}(E)}_{F(o)}$:

$$\langle R'_g(H), R_{\tilde{g}}(f) \rangle_{\mathcal{D}'_{F(o)}(E)} = \langle H, f \rangle_{\mathcal{D}_{F(o)}(E)}. \quad (126)$$

From formulas (126), (122), and (102) we obtain the following formula for any $f \in \mathcal{E}(E)$, any $H \in \mathcal{D}'_{F(o)}(E)$, and any $\tilde{g} \in \widehat{\text{Aut}_{C_2}(E)}_{F(o)}$:

$$R'_{\tilde{g}}(f \cdot H) = R_{\Lambda(\tilde{g})}(f) \cdot R'_{\tilde{g}}(H).$$

5.5.5

Let $E = (I, F, V)$ be a complete C_2 -space over the field \mathbb{F}_q . Then $\check{E} = (I^0, F^0, \check{V})$, and $\check{\check{E}} = E$. For any $g \in \text{Aut}_{C_2}(E)$ we have canonically $\check{g} \in \text{Aut}_{C_2}(\check{E})$. And we have the following isomorphism of groups:

$$g \in \text{Aut}_{C_2}(E) \longmapsto \check{g}^{-1} \in \text{Aut}_{C_2}(\check{E}). \quad (127)$$

Also for any $o \in I$ we have the following isomorphism of groups:

$$(g, \mu) \in \widehat{\text{Aut}_{C_2}(E)}_{F(o)} \longmapsto (\check{g}^{-1}, \mu) \in \widehat{\text{Aut}_{C_2}(\check{E})}_{F^0(o)},$$

where we use isomorphism (103) to obtain that for any $g \in \text{Aut}_{C_2}(E)$ canonically

$$\mu(F(o) \mid gF(o)) = \mu(F^0(o) \mid \check{g}^{-1}F^0(o)). \quad (128)$$

Using isomorphism (127) we obtain representations of the group $\text{Aut}_{C_2}(E)$ on \mathbf{C} -spaces $\mathcal{E}(\check{E})$, $\tilde{\mathcal{E}}(\check{E})$, $\mathcal{E}'(\check{E})$, $\tilde{\mathcal{E}}'(\check{E})$.

Using isomorphism (128) we obtain representations of the group $\widehat{\text{Aut}_{C_2}(E)}_{F(o)}$ on \mathbf{C} -spaces $\mathcal{D}_{F^0(o)}(\check{E})$ and $\mathcal{D}'_{F^0(o)}(\check{E})$.

We have the following proposition.

Proposition 25 *Let $E = (I, F, V)$ be a complete C_2 -space over the field \mathbb{F}_q . We fix some $o \in I$. Then*

1. *The Fourier transform \mathbf{F} gives an isomorphism between representations of the group $\text{Aut}_{C_2}(E)$ on \mathbf{C} -spaces $\mathcal{E}(E)$ and $\tilde{\mathcal{E}}'(\check{E})$ and on \mathbf{C} -spaces $\tilde{\mathcal{E}}(E)$ and $\mathcal{E}'(\check{E})$.*
2. *The Fourier transform \mathbf{F} gives an isomorphism between representations of the group $\widehat{\text{Aut}_{C_2}(E)}_{F(o)}$ on \mathbf{C} -spaces $\mathcal{D}_{F(o)}(E)$ and $\mathcal{D}_{F^0(o)}(\check{E})$ and on \mathbf{C} -spaces $\mathcal{D}'_{F(o)}(E)$ and $\mathcal{D}'_{F^0(o)}(\check{E})$.*

Proof . Using definition of 2-dimensional Fourier transform \mathbf{F} from section 5.4.2, and using formulas (93)-(98), we reduce the statements of this proposition to corresponding statements about isomorphisms of C_1 -spaces and 1-dimensional Fourier transforms between them. The last statements follow from the definition and properties of 1-dimensional Fourier transform, or can be again reduced to 0-dimensional case by formulas (10)-(15) using remark 6. The corresponding statement in the 0-dimensional case looks as following. For any finite-dimensional vector space W over the field \mathbb{F}_q , for any $f \in \mathcal{F}(W)$, for any $u \in W^*$, for any $g \in \text{Aut}_{\mathbb{F}_q}(W)$ we have:

$$\sum_{w \in W} f(g^{-1}w) \overline{\psi(u(w))} = \sum_{w \in W} f(w) \overline{\psi(u(gw))}.$$

5.6 Direct and inverse images

In this section we construct direct and inverse images of spaces of functions and distributions on C_2 -spaces.

5.6.1

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_2 -spaces over a finite field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. By definition, there are order-preserving functions:

- (i) $\gamma : I_2 \longrightarrow I_3$ such that $\beta(F_2(i)) = F_3(\gamma(i))$ for any $i \in I_2$,
- (ii) $\varepsilon : I_2 \longrightarrow I_1$ such that $F_2(i) \cap V_1 = F_1(\varepsilon(i))$ for any $i \in I_2$.

Proposition 26 *We suppose that E_1 is a cC_2 -space. Let $o \in I_2$. Then there is the direct image*

$$\beta_* : \mathcal{D}_{F_2(o)}(E_2) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(o)) \mid V_1) \longrightarrow \mathcal{D}_{F_3(\gamma(o))}(E_3).$$

Proof . For any $i, j \in I_2$ we have canonically

$$\mu(F_2(i) \mid F_2(j)) = \mu(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))) \otimes_{\mathbf{C}} \mu(F_3(\gamma(i)) \mid F_3(\gamma(j))). \quad (129)$$

Let $k \in I_2$ such that $F_1(\varepsilon(k)) = V_1$. Let $i \geq j \in I_2$ be any such that $i \geq k \geq j$. Then we have an admissible triple of C_1 -spaces:

$$0 \longrightarrow V_1/F_1(\varepsilon(j)) \longrightarrow F_2(i)/F_2(j) \xrightarrow{\beta_{ij}} F_3(\gamma(i))/F_3(\gamma(j)) \longrightarrow 0.$$

We have a well-defined map:

$$(\beta_{ij})_* : \mathcal{D}(F_2(i)/F_2(j)) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(j)) \mid V_1) \longrightarrow \mathcal{D}(F_3(\gamma(i))/F_3(\gamma(j))),$$

where we use that $\mu(F_1(\varepsilon(j)) \mid V_1) = \mu(V_1/F_1(\varepsilon(j)))$.

Therefore we have a well-defined map

$$\begin{aligned} & (\beta_{ij})_* \otimes \mu(F_3(\gamma(j)) \mid F_3(\gamma(o))) : \\ & \mathcal{D}(F_2(i)/F_2(j)) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(j)) \mid V_1) \otimes_{\mathbf{C}} \mu(F_3(\gamma(j)) \mid F_3(\gamma(o))) \longrightarrow \\ & \longrightarrow \mathcal{D}(F_3(\gamma(i)) \mid F_3(\gamma(j))) \otimes_{\mathbf{C}} \mu(F_3(\gamma(j)) \mid F_3(\gamma(o))). \end{aligned}$$

From (129) we have that

$$\mu(F_3(\gamma(j)) \mid F_3(\gamma(o))) = \mu(F_2(j) \mid F_2(o)) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(o)) \mid F_1(\varepsilon(j))).$$

Therefore

$$\mu(F_1(\varepsilon(j)) \mid V_1) \otimes_{\mathbf{C}} \mu(F_3(\gamma(j)) \mid F_3(\gamma(o))) = \mu(F_2(j) \mid F_2(o)) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(o)) \mid V_1).$$

Thus we have a well-defined map

$$\begin{aligned} & (\beta_{ij})_* \otimes \mu(F_3(\gamma(j)) \mid F_3(\gamma(o))) : \\ & \mathcal{D}(F_2(i)/F_2(j)) \otimes_{\mathbf{C}} \mu(F_2(j) \mid F_2(o)) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(o)) \mid V_1) \longrightarrow \\ & \longrightarrow \mathcal{D}(F_3(\gamma(i))/F_3(\gamma(j))) \otimes_{\mathbf{C}} \mu(F_3(\gamma(j)) \mid F_3(\gamma(o))). \end{aligned}$$

Now we take the projective limits which are used to construct spaces $\mathcal{D}_{F_2(o)}(E_2)$ and $\mathcal{D}_{F_3(\gamma(o))}(E_3)$. Then we obtain a well-defined map β_* .

Remark 16 Dually to the maps above, we obtain a well-defined map

$$\beta^* : \mathcal{D}'_{F_3(\gamma(o))}(E_3) \otimes_{\mathbf{C}} \mu(F_3(\varepsilon(o)) \mid V_1) \longrightarrow \mathcal{D}'_{F_2(o)}(E_2)$$

such that the maps β_* and β^* are conjugate maps with respect to the pairings $\langle \cdot, \cdot \rangle_{\mathcal{D}_{F_2(o)}(E_2)}$ and $\langle \cdot, \cdot \rangle_{\mathcal{D}_{F_3(\gamma(o))}(E_3)}$.

5.6.2

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_2 -spaces over a finite field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. By definition, there are order-preserving functions:

- (i) $\gamma : I_2 \longrightarrow I_3$ such that $\beta(F_2(i)) = F_3(\gamma(i))$ for any $i \in I_2$,

(ii) $\varepsilon : I_2 \rightarrow I_1$ such that $F_2(i) \cap V_1 = F_1(\varepsilon(i))$ for any $i \in I_2$.

Proposition 27 *We suppose that E_3 is a dC_2 -space. Let $o \in I_2$. Then there is the inverse image*

$$\alpha^* : \mathcal{D}_{F_2(o)}(E_2) \otimes_{\mathbf{C}} \mu(F_3(\gamma(o)) \mid \{0\}) \rightarrow \mathcal{D}_{F_1(\varepsilon(o))}(E_1),$$

where $\{0\}$ is the zero subspace of V_3 .

Proof . For any $i, j \in I_2$ we have canonically

$$\mu(F_2(i) \mid F_2(j)) = \mu(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))) \otimes_{\mathbf{C}} \mu(F_3(\gamma(i)) \mid F_3(\gamma(j))). \quad (130)$$

Since β is an admissible epimorphism and E_3 is a dC_2 -space, there is $k \in I_2$ such that $F_3(\gamma(k)) = \{0\}$. Let $i \geq j \in I_2$ be any such that $i \geq k \geq j$. Then we have an admissible triple of C_1 -spaces:

$$0 \rightarrow F_1(\varepsilon(i))/F_1(\varepsilon(j)) \xrightarrow{\alpha_{ij}} F_2(i)/F_2(j) \rightarrow F_3(\gamma(i)) \rightarrow 0. \quad (131)$$

We have a well-defined map

$$\alpha_{ij}^* : \mathcal{D}(F_2(i)/F_2(j)) \rightarrow \mathcal{D}(F_1(\varepsilon(i))/F_1(\varepsilon(j))).$$

Therefore we have a well-defined map

$$\begin{aligned} & \alpha_{ij}^* \otimes \mu(F_1(\varepsilon(j)) \mid F_1(\varepsilon(o))) : \\ & \mathcal{D}(F_2(i)/F_2(j)) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(j)) \mid F_1(\varepsilon(o))) \rightarrow \\ & \rightarrow \mathcal{D}(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(j)) \mid F_1(\varepsilon(o))). \end{aligned}$$

From (130) and (131) we have that

$$\mu(F_2(o) \mid F_2(j)) = \mu(F_1(\varepsilon(o)) \mid F_1(\varepsilon(j))) \otimes_{\mathbf{C}} \mu(F_3(\gamma(o)) \mid \{0\}).$$

Hence

$$\mu(F_1(\varepsilon(j)) \mid F_1(\varepsilon(o))) = \mu(F_2(j) \mid F_2(o)) \otimes_{\mathbf{C}} \mu(F_3(\gamma(o)) \mid \{0\}).$$

Thus we have a well-defined map

$$\begin{aligned} & \alpha_{ij}^* \otimes \mu(F_1(\varepsilon(j)) \mid F_1(\varepsilon(o))) : \\ & \mathcal{D}(F_2(i)/F_2(j)) \otimes_{\mathbf{C}} \mu(F_2(j)/F_2(o)) \otimes_{\mathbf{C}} \mu(F_3(\gamma(o)) \mid \{0\}) \rightarrow \\ & \rightarrow \mathcal{D}(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(j)) \mid F_1(\varepsilon(o))). \end{aligned}$$

Now we take the projective limits (with respect to $i \geq k \geq j$) which we used to construct spaces $\mathcal{D}_{F_2(o)}(E_2)$ and $\mathcal{D}_{F_1(\varepsilon(o))}(E_1)$. Then we obtain a well-defined map α^* .

Remark 17 Dually to the maps above, we obtain a well-defined map

$$\alpha_* : \mathcal{D}'_{F_1(\varepsilon(o))}(E_1) \otimes_{\mathbf{C}} \mu(F_3(\gamma(o)) \mid \{0\}) \rightarrow \mathcal{D}'_{F_2(o)}(E_2)$$

such that the maps α^* and α_* are conjugate maps with respect to the pairings $\langle \cdot, \cdot \rangle_{\mathcal{D}_{F_2(o)}(E_2)}$ and $\langle \cdot, \cdot \rangle_{\mathcal{D}_{F_1(\varepsilon(o))}(E_1)}$.

5.6.3

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_2 -spaces over a finite field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. By definition, there are order-preserving functions:

- (i) $\gamma : I_2 \longrightarrow I_3$ such that $\beta(F_2(i)) = F_3(\gamma(i))$ for any $i \in I_2$,
- (ii) $\varepsilon : I_2 \longrightarrow I_1$ such that $F_2(i) \cap V_1 = F_1(\varepsilon(i))$ for any $i \in I_2$.

Proposition 28 *We suppose that E_1 is a cfC_2 -space. Let $o \in I_2$. Then there is the inverse image*

$$\beta^* : \mathcal{D}_{F_3(\gamma(o))}(E_3) \longrightarrow \mathcal{D}_{F_2(o)}(E_2).$$

Proof . For any $i, j \in I_2$ we have canonically

$$\mu(F_2(i) \mid F_2(j)) = \mu(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))) \otimes_{\mathbf{C}} \mu(F_3(\gamma(i)) \mid F_3(\gamma(j))). \quad (132)$$

Since E_1 is a cfC_2 -space, for any $i \geq j \in I_2$ we have a canonical element $1_{ij} \in \mu(F_1(\varepsilon(i))/F_1(\varepsilon(j)))$ such that $1_{ij}(\delta_{F_1(\varepsilon(i))|F_1(\varepsilon(j))}) = 1$.

Therefore we have a canonical element $1_{ij} \in \mu(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j)))$ for any $i, j \in I_2$ such that $1_{ij} \otimes 1_{jk} = 1_{ik}$ for any $i, j, k \in I_2$. Thus, we have canonically that $\mu(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))) = \mathbf{C}$ for any $i, j \in I_2$. (See also remark 12).

Thus from formula (132) we have for any $i, j \in I_2$

$$\mu(F_2(i) \mid F_2(j)) = \mu(F_3(\gamma(i)) \mid F_3(\gamma(j))). \quad (133)$$

For any $i \geq j \in I_2$ we have an admissible triple of C_1 -spaces:

$$0 \longrightarrow \frac{F_1(\varepsilon(i))}{F_1(\varepsilon(j))} \longrightarrow \frac{F_2(i)}{F_2(j)} \xrightarrow{\beta_{ij}} \frac{F_3(\gamma(i))}{F_3(\gamma(j))} \longrightarrow 0.$$

We have a well-defined map:

$$\beta_{ij}^* : \mathcal{D}(F_3(\gamma(i))/F_3(\gamma(j))) \longrightarrow \mathcal{D}(F_2(i)/F_2(j)).$$

From formula (133) we have that

$$\mu(F_2(j) \mid F_2(o)) = \mu(F_3(\gamma(j)) \mid F_3(\gamma(o))).$$

Therefore we have a map:

$$\begin{aligned} \beta_{ij}^* \otimes \mu(F_2(j) \mid F_2(o)) & : \\ \mathcal{D}(F_3(\gamma(i))/F_3(\gamma(j))) \otimes_{\mathbf{C}} \mu(F_3(\gamma(j)) \mid F_3(\gamma(o))) & \longrightarrow \\ & \longrightarrow \mathcal{D}(F_2(i)/F_2(j)) \otimes_{\mathbf{C}} \mu(F_2(j) \mid F_2(o)). \end{aligned}$$

These maps (for various $i \geq j \in I_2$) are compatible when we take the projective limits according to formulas which define $\mathcal{D}_{F_3(\gamma(o))}(E_3)$ and $\mathcal{D}_{F_2(o)}(E_2)$. Therefore we obtain the map β^* .

Remark 18 Dually to the maps, which we considered above, we have a well-defined map

$$\beta_* : \mathcal{D}'_{F_2(o)}(E_2) \longrightarrow \mathcal{D}'_{F_3(\gamma(o))}(E_3)$$

such that the maps β^* and β_* are conjugate maps with respect to the pairings $\langle \cdot, \cdot \rangle_{\mathcal{D}_{F_2(o)}(E_2)}$ and $\langle \cdot, \cdot \rangle_{\mathcal{D}_{F_3(\gamma(o))}(E_3)}$.

5.6.4

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_2 -spaces over a finite field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. By definition, there are order-preserving functions:

- (i) $\gamma : I_2 \longrightarrow I_3$ such that $\beta(F_2(i)) = F_3(\gamma(i))$ for any $i \in I_2$,
- (ii) $\varepsilon : I_2 \longrightarrow I_1$ such that $F_2(i) \cap V_1 = F_1(\varepsilon(i))$ for any $i \in I_2$.

Proposition 29 *We suppose that E_3 is a dfC_2 space. Let $o \in I_2$. Then there is the direct image*

$$\alpha_* : \mathcal{D}_{F_1(\varepsilon(o))}(E_1) \longrightarrow \mathcal{D}_{F_2(o)}(E_2).$$

Proof . For any $i, j \in I_2$ we have canonically

$$\mu(F_2(i) \mid F_2(j)) = \mu(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))) \otimes_{\mathbf{C}} \mu(F_3(\gamma(i)) \mid F_3(\gamma(j))). \quad (134)$$

Since E_3 is a dfC_2 -space, for any $i \geq j \in I_2$ we have a canonical element $\delta_{ij} \in \mu(F_3(\gamma(i))/F_3(\gamma(j)))$ such that $\delta_{ij}(\{0\}) = 1$. Therefore we have a canonical element $\delta_{ij} \in \mu(F_3(\gamma(i)) \mid F_3(\gamma(j)))$ for any $i, j \in I_2$ such that $\delta_{ij} \otimes \delta_{jk} = \delta_{ik}$ for any $i, j, k \in I_2$. Therefore we have canonically $\mu(F_3(\gamma(i)) \mid F_3(\gamma(j))) = \mathbf{C}$ for any $i, j \in I_2$. (See also remark 12).

Thus from formula (134) we have for any $i, j \in I_2$

$$\mu(F_2(i) \mid F_2(j)) = \mu(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))). \quad (135)$$

For any $i \geq j \in I_2$ we have an admissible triple of C_1 -spaces:

$$0 \longrightarrow \frac{F_1(\varepsilon(i))}{F_1(\varepsilon(j))} \xrightarrow{\alpha_{ij}} \frac{F_2(i)}{F_2(j)} \longrightarrow \frac{F_3(\gamma(i))}{F_3(\gamma(j))} \longrightarrow 0.$$

We have a well-defined map:

$$(\alpha_{ij})_* : \mathcal{D}(F_1(\varepsilon(i))/F_1(\varepsilon(j))) \longrightarrow \mathcal{D}(F_2(i)/F_2(j)),$$

because $\frac{F_3(\gamma(i))}{F_3(\gamma(j))}$ is a discrete C_1 -space.

From formula (135) we have that

$$\mu(F_2(j) \mid F_2(o)) = \mu(F_1(\varepsilon(j)) \mid F_1(\varepsilon(o))).$$

Therefore we have a map:

$$\begin{aligned} & (\alpha_{ij})_* \otimes \mu(F_2(j) \mid F_2(o)) : \\ & \mathcal{D}(F_1(\varepsilon(i))/F_1(\varepsilon(j))) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(j)) \mid F_1(\varepsilon(o))) \longrightarrow \\ & \longrightarrow \mathcal{D}(F_2(i)/F_2(j)) \otimes_{\mathbf{C}} \mu(F_2(j) \mid F_2(o)). \end{aligned}$$

These maps (for various $i \geq j \in I_2$) are compatible when we take the projective limits according to the formulas which define $\mathcal{D}_{F_1(\varepsilon(o))}(E_1)$ and $\mathcal{D}_{F_2(o)}(E_2)$. Therefore we obtain the map α_* .

Remark 19 Dually to the maps, which we considered above, we have a well-defined map

$$\alpha^* : \mathcal{D}'_{F_2(o)}(E_2) \longrightarrow \mathcal{D}'_{F_1(\varepsilon(o))}(E_1)$$

such that the maps α_* and α^* are conjugate maps with respect to the pairings $\langle \cdot, \cdot \rangle_{\mathcal{D}_{F_2(o)}(E_2)}$ and $\langle \cdot, \cdot \rangle_{\mathcal{D}_{F_1(\varepsilon(o))}(E_1)}$.

5.7 Composition of maps and base change rules.

In this section we consider base change rules (subsection 5.7.1) and rules of composition of maps (subsection 5.7.2).

5.7.1

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_2 -spaces over the field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$.

Let

$$0 \longrightarrow D \xrightarrow{\gamma} E_3 \xrightarrow{\delta} B \longrightarrow 0$$

be an admissible triple of C_2 -spaces over the field \mathbb{F}_q , where $D = (R, S, Y)$, $B = (T, U, W)$. Then we have the following commutative diagram of C_2 -spaces.

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & E_1 & \xrightarrow{\gamma\alpha} & E_2 \times_{E_3} D & \xrightarrow{\gamma\beta} & D \longrightarrow 0 \\
 & & \parallel & & \downarrow \beta_\gamma & & \downarrow \gamma \\
 0 & \longrightarrow & E_1 & \xrightarrow{\alpha} & E_2 & \xrightarrow{\beta} & E_3 \longrightarrow 0 \\
 & & & & \downarrow \beta_\delta & & \downarrow \delta \\
 & & & & B & \xlongequal{\quad} & B \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array} \tag{136}$$

Let $X' = E_2 \times_{E_3} D = (N, Q, X)$ as C_2 -space. In this diagram two horizontal triples and two vertical triples are admissible triples of C_2 -spaces.

We choose any $i \leq j \in I_2$. Then diagram (136) induces the following commutative diagram of C_1 -spaces (see the construction of fibered product from lemma 5).

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \frac{F_2(j) \cap V_1}{F_2(i) \cap V_1} & \xrightarrow{(\gamma_\alpha)_{ji}} & \frac{F_2(j) \cap X}{F_2(i) \cap X} & \xrightarrow{(\gamma_\beta)_{ji}} & \frac{\beta(F_2(j)) \cap Y}{\beta(F_2(i)) \cap Y} \longrightarrow 0 \\
& & \parallel & & \downarrow (\beta_\gamma)_{ji} & & \downarrow \gamma_{ji} \\
0 & \longrightarrow & \frac{F_2(j) \cap V_1}{F_2(i) \cap V_1} & \xrightarrow{\alpha_{ji}} & \frac{F_2(j)}{F_2(i)} & \xrightarrow{\beta_{ji}} & \frac{\beta(F_2(j))}{\beta(F_2(i))} \longrightarrow 0 \\
& & & & \downarrow (\beta_\delta)_{ji} & & \downarrow \delta_{ji} \\
& & & & \frac{\delta\beta(F_2(j))}{\delta\beta(F_2(i))} & = & \frac{\delta\beta(F_2(j))}{\delta\beta(F_2(i))} \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0
\end{array}
\tag{137}$$

Here two horizontal triples and two vertical triples are admissible triples of C_1 -spaces. At the same time,

$$\frac{F_2(j) \cap X}{F_2(i) \cap X} = \frac{F_2(j)}{F_2(i)} \times_{\frac{\beta(F_2(j))}{\beta(F_2(i))}} \frac{\beta(F_2(j)) \cap Y}{\beta(F_2(i)) \cap Y}$$

as C_1 -spaces.

Proposition 30 . Using notations of diagram (136), let E_1 be a cC_2 -space, B be a dC_2 -space. Let $o \in I_2$, $\mu \in \mu(F_2(o) \cap V_1 \mid V_1)$, $\nu \in \mu(\delta\beta(F_2(o)) \mid \{0\})$. Then the following formulas are satisfied.

1. For any $f \in \mathcal{D}_{F_2(o)}(E_2)$

$$\gamma^*(\beta_*(f \otimes \mu) \otimes \nu) = (\gamma_\beta)_*(\beta_\gamma^*(f \otimes \nu) \otimes \mu). \tag{138}$$

2. For any $G \in \mathcal{D}'_{\beta(F_2(o)) \cap Y}(D)$

$$\beta^*(\gamma_*(G \otimes \nu) \otimes \mu) = (\beta_\gamma)_*(\gamma_\beta^*(G \otimes \mu) \otimes \nu). \tag{139}$$

Remark 20 . The formulation of this proposition is well-defined (with respect to definitions of direct and inverse images), since $\alpha = \beta_\gamma \circ \gamma_\alpha$, $\beta_\delta = \delta \circ \beta$.

Proof of proposition 30. We prove at first formula (138). Let $j_0 \in I_2$ be such that $F_2(j_0) \cap V_1 = V_1$. (Such a j_0 it is possible to choose, since E_1 is a cC_2 -space.) Let $i_0 \in I_2$ be such that $\beta\delta(F_2(i_0)) = \{0\}$. (Such an i_0 it is possible to choose, since B is a dC_2 -space.) We consider arbitrary $j \geq i \in I_2$ such that $j \geq j_0$, $i \leq i_0$. Then the base change formula on C_1 -spaces is satisfied for the maps β_{ji} , γ_{ji} , $(\gamma_\beta)_{ji}$, $(\beta_\gamma)_{ji}$ from cartesian square of diagram (137) (see formula (60) from proposition 18). We multiply this formula by corresponding measure spaces, and then we take the projective limit with respect to all such $j \geq i \in I_2$ ($j \geq j_0$, $i \leq i_0$). Using the constructions of maps β_* and γ^* from propositions 26 and 27, we obtain the base change formula (138).

Formula (139) is a dual formula to formula (138), and it can be obtained by analogous reasonings as in the proof of formula (138), but one has to take the inductive limit with respect to $j, i \in I_2$, and one has to use formula (61) instead of formula (60).

Proposition 31 . Using notations of diagram (136), let E_1 be a cfC_2 -space, B be a dfC_2 -space. Let $o \in I_2$. Then the following formulas are satisfied.

1. For any $f \in \mathcal{D}_{\beta(F_2(o)) \cap Y}(D)$

$$\beta^* \gamma_*(f) = (\beta_\gamma)_* \gamma_\beta^*(f). \quad (140)$$

2. For any $G \in \mathcal{D}'_{F_2(o)}(E_2)$

$$\gamma^* \beta_*(G) = (\gamma_\beta)_* \beta_\gamma^*(G). \quad (141)$$

Proof . We prove formula (140). We consider arbitrary $j \geq i \in I_2$. Then the base change formula on C_1 -spaces is satisfied for the maps β_{ji} , γ_{ji} , $(\gamma_\beta)_{ji}$, $(\beta_\gamma)_{ji}$ from cartesian square of diagram (137) (see formula (68) from proposition 18). We take the projective limit with respect to such $j \geq i \in I_2$. Using the constructions of maps β^* and γ_* from propositions 28 and 29, we obtain the base change formula (140).

Formula (141) can be obtained by analogous reasonings, but one has to take the inductive limit with respect to $i, j \in I_2$, and one has to use formula (69) instead of formula (68).

Proposition 32 . Using notations of diagram (136), let E_1 be a cC_2 -space, B be a dfC_2 -space. Let $o \in I_2$, $\mu \in \mu(F_2(o) \cap V_1 \mid V_1)$. Then the following formulas are satisfied.

1. For any $f \in \mathcal{D}_{F_2(o) \cap X}(X')$, $\mu \in \mu(E_1)$

$$\beta_*((\beta_\gamma)_*(f) \otimes \mu) = \gamma_*(\gamma_\beta)_*(f \otimes \mu). \quad (142)$$

2. For any $G \in \mathcal{D}'_{\beta(F_2(o))}(E_3)$, $\mu \in \mu(E_1)$

$$\gamma_\beta^*(\gamma^*(G) \otimes \mu) = \beta_\gamma^* \beta^*(G \otimes \mu). \quad (143)$$

Proof is analogous to the proof of proposition 30 and 31, and it is reduced to corresponding formula (66) for C_1 -spaces (to prove formula (142)) and to corresponding formula (67) for C_1 -spaces (to prove formula (143)).

Proposition 33 . Using notations of diagram (136), let E_1 be a cC_2 -space, B be a dC_2 -space. Let $o \in I_2$, $\nu \in \mu(\delta\beta(F_2(o)) \mid \{0\})$. Then the following formulas are satisfied.

1. For any $f \in \mathcal{D}_{\beta(F_2(o))}(E_3)$

$$\gamma_\beta^* \gamma^*(f \otimes \nu) = \beta_\gamma^*(\beta^*(f) \otimes \nu). \quad (144)$$

2. For any $G \in \mathcal{D}'_{F_2(o) \cap X}(X')$

$$\beta_*(\beta_\gamma)_*(G \otimes \nu) = \gamma_*((\gamma_\beta)_*(G) \otimes \nu). \quad (145)$$

Proof is analogous to the proof of proposition 30 and 31, and it is reduced to corresponding formula (64) for C_1 -spaces (to prove formula (144)) and to corresponding formula (65) for C_1 -spaces (to prove formula (145)).

5.7.2

We consider again diagram (136). If E_1 and D are cC_2 -spaces, then $E_2 \times_{E_3} D$ is a cC_2 -space. Indeed, it follows from the admissible triple of C_2 -spaces:

$$0 \longrightarrow E_1 \xrightarrow{\gamma_\alpha} E_2 \times_{E_3} D \xrightarrow{\gamma_\beta} D \longrightarrow 0. \quad (146)$$

Let $o \in I_2$. Then from (146) we obtain the canonical isomorphism

$$\mu(F_2(o) \cap X \mid X) = \mu(F_2(o) \cap V_1 \mid V_1) \otimes_{\mathbf{C}} \mu(\beta(F_2(o)) \cap Y \mid Y). \quad (147)$$

(The subspaces, which appear in formula (147), are elements of the filtration of corresponding C_2 -spaces. Therefore the spaces of virtual measures are well-defined, see section 5.2.)

We have from diagram (136) the following admissible triple of C_2 -spaces:

$$0 \longrightarrow E_2 \times_{E_3} D \xrightarrow{\beta_\gamma} E_2 \xrightarrow{\delta\beta} B \longrightarrow 0.$$

Proposition 34 . Using notations of diagram (136), let E_1 and D be cC_2 -spaces. Let $o \in I_2$, $\mu \in \mu(F_2(o) \cap V_1 \mid V_1)$, $\nu \in \mu(\beta(F_2(o)) \cap Y \mid Y)$. Then the following formulas are satisfied.

1. For any $f \in \mathcal{D}_{F_2(o)}(E_2)$

$$(\delta\beta)_*(f \otimes (\mu \otimes \nu)) = \delta_*(\beta_*(f \otimes \mu) \otimes \nu). \quad (148)$$

2. For any $G \in \mathcal{D}'_{\delta\beta(F_2(o))}(B)$

$$(\delta\beta)^*(G \otimes (\mu \otimes \nu)) = \beta^*(\delta^*(G \otimes \nu) \otimes \mu). \quad (149)$$

Remark 21 . Due to (147) we have that $\mu \otimes \nu \in \mu(F_2(o) \cap X \mid X)$.

Proof of proposition 34. Let $j_0 \in I_2$ be such that $F_2(j_0) \cap V_1 = V_1$ and $\beta(F_2(j_0)) \cap Y = Y$. We consider arbitrary $j \geq i \in I_2$ such that $j \geq j_0$. Then formula (43) from proposition 16 is satisfied for the maps β_{ji} , δ_{ji} and $(\beta_\delta)_{ji} = \delta_{ji}\beta_{ji}$ (see diagram (137)). We multiply this formula by corresponding measure spaces, and then we take the limit with respect to $j \geq i \in I_2$ ($j \geq j_0$). Using explicit construction of direct image from proposition 26, we obtain formula (148).

Formula (149) can be obtained by analogous reasonings, but one has to use formula (44) from proposition 16.

Using notations of diagram (136), we suppose that E_1 and D are cfC_2 -spaces. Then from admissible triple of C_2 -spaces (146) it follows that $E_2 \times_{E_3} D$ is also a cfC_2 -space.

Proposition 35 . Using notations of diagram (136), let E_1 and D be cfC_2 -spaces. Let $o \in I_2$. Then the following formulas are satisfied.

1. For any $f \in \mathcal{D}_{\delta\beta(F_2(o))}(B)$

$$(\delta\beta)^*(f) = \beta^*\delta^*(f). \quad (150)$$

2. For any $G \in \mathcal{D}'_{F_2(o)}(E_2)$

$$(\delta\beta)_*(G) = \delta_*\beta_*(G). \quad (151)$$

Proof . We consider arbitrary $j \geq i \in I_2$. Then formula (47) from proposition 16 is satisfied for the maps β_{ji} , δ_{ji} and $(\beta_\delta)_{ji} = \delta_{ji}\beta_{ji}$ (see diagram (137)). We multiply this formula by corresponding measure spaces, and then we take the limit with respect to $j \geq i \in I_2$. Using explicit construction of inverse image from proposition 28, we obtain formula (150).

Formula (151) can be obtained by analogous reasonings, but one has to use formula (48) from proposition 16.

A little bit rewriting (and redenoting) diagram (136), we obtain the following

commutative diagram of C_2 -spaces over the field \mathbb{F}_q :

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & E_1 & \xrightarrow{\alpha} & E_2 & \xrightarrow{\beta} & E_3 \longrightarrow 0 \\
& & \parallel & & \downarrow \alpha' & & \downarrow \beta_{\alpha'} \\
0 & \longrightarrow & E_1 & \xrightarrow{\alpha' \alpha} & H' & \xrightarrow{\alpha'_{\beta}} & E_3 \amalg_{E_2} H' \longrightarrow 0 \\
& & & & \downarrow \theta & & \downarrow \theta' \\
& & & & L' & \xlongequal{\quad} & L' \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0
\end{array} \tag{152}$$

In this diagram two horizontal and two vertical triples are admissible triples of C_2 -spaces.

Let $E_i = (I_i, F_i, V_i)$ ($1 \leq i \leq 3$), $H' = (J', T', W')$ as C_2 -spaces. From admissible triple of C_2 -spaces

$$0 \longrightarrow E_3 \xrightarrow{\beta_{\alpha'}} E_3 \amalg_{E_2} H' \xrightarrow{\theta'} L' \longrightarrow 0 \tag{153}$$

we obtain that if E_3 and L' are dC_2 -spaces, then $E_3 \amalg_{E_2} H'$ is also a dC_2 -space.

Let $o \in J'$ be any, then from (153) we canonically have

$$\mu(\alpha'_{\beta}(T'(o)) \mid \{0\}) = \mu(\beta(T'(o) \cap V_2) \mid \{0\}) \otimes_{\mathbf{C}} \mu(\theta(T'(o)) \mid \{0\}). \tag{154}$$

(The subspaces, which appear in formula (154), are elements of the filtration of corresponding C_2 -spaces. Therefore the spaces of virtual measures are well-defined, see section 5.2.)

From diagram (152) we have an admissible triple

$$0 \longrightarrow E_1 \xrightarrow{\alpha' \alpha} H' \xrightarrow{\alpha'_{\beta}} E_3 \amalg_{E_2} H' \longrightarrow 0.$$

Proposition 36 . Using notations of diagram (152), let E_3 and L' be dC_2 -spaces. Let $o \in J'$, $\mu \in \mu(\beta(T'(o) \cap V_2) \mid \{0\})$, $\nu \in \mu(\theta(T'(o)) \mid \{0\})$. Then the following formulas are satisfied.

1. For any $f \in \mathcal{D}_{T'(o)}(H')$

$$(\alpha' \alpha)^*(f \otimes (\mu \otimes \nu)) = \alpha^*((\alpha')^*(f \otimes \nu) \otimes \mu). \quad (155)$$

2. For any $G \in \mathcal{D}'_{T'(o) \cap V_1}(E_1)$

$$(\alpha' \alpha)_*(G \otimes (\mu \otimes \nu)) = (\alpha')_*(\alpha_*(G \otimes \mu) \otimes \nu). \quad (156)$$

Remark 22 . Due to (154) we have that $\mu \otimes \nu \in \mu(\alpha'_\beta(T'(o)) \mid \{0\})$.

Proof of proposition. Formula (155) can be reduced to formula (52) from proposition 17 after the taking the projective limit with respect to $j \geq i \in J'$, where $i \leq i_0$, and i_0 is chosen such that $\theta(T'(i_0)) = \{0\}$ and $\beta(T'(i_0) \cap V_2) = \{0\}$. At that we use the explicit construction of inverse image from proposition 27. (Compare with the proof of proposition 35.)

Formula (156) can be proved analogously, but one has to use formula (53) from proposition 17 and take the inductive limit.

Using notations of diagram (152) we suppose that E_3 and L' are dfC_2 -spaces. Then from admissible triple of C_2 -spaces (153) it follows that $E_3 \amalg_{E_2} H'$ is also a dfC_2 -space.

Proposition 37 . Using notations of diagram (152), let E_3 and L' are dfC_2 -spaces. Let $o \in J'$. Then the following formulas are satisfied.

1. For any $f \in \mathcal{D}_{T'(o) \cap V_1}(E_1)$

$$(\alpha' \alpha)_*(f) = (\alpha')_* \alpha_*(f). \quad (157)$$

2. For any $G \in \mathcal{D}'_{T'(o)}(H')$

$$(\alpha' \alpha)^*(G) = \alpha^*((\alpha')^*(G)). \quad (158)$$

Proof of formula (157) can be reduced to formula (56) of proposition 17 after the choice of $j \geq i \in J'$ and taking the projective limit with respect to this $j \geq i \in J'$. At that we use the explicit construction of direct image from proposition 29.

The proof of formula (158) is analogous, but one has to take the inductive limit with respect to $j \geq i \in J'$ and use formula (57) instead of formula (56).

5.8 Fourier transform and direct and inverse images

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of C_2 -spaces over a finite field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. We suppose that E_i ($1 \leq i \leq 3$) is a *complete* C_2 -space. Then $\bar{E}_i = E_i$, $1 \leq i \leq 3$.

By definition, there are order-preserving functions:

(i) $\gamma: I_2 \longrightarrow I_3$ such that $\beta(F_2(i)) = F_3(\gamma(i))$ for any $i \in I_2$;

(ii) $\varepsilon : I_2 \longrightarrow I_1$ such that $F_2(i) \cap V_1 = F_1(\varepsilon(i))$ for any $i \in I_2$.

We consider the following admissible triple:

$$0 \longrightarrow \check{E}_3 \xrightarrow{\check{\beta}} \check{E}_2 \xrightarrow{\check{\alpha}} \check{E}_1 \longrightarrow 0.$$

We have $\check{E}_i = (I_i^0, F_i^0, \check{V}_i)$, $1 \leq i \leq 3$. We recall that the partially ordered set I_i^0 ($1 \leq i \leq 3$) is equal to I_i as a set, but has inverse partial order comparing with I_i ($1 \leq i \leq 3$). We have that for any $i, j \in I_2$:

$$\begin{aligned} \check{\alpha}(F_2^0(i)) &= F_1^0(\varepsilon(i)) \\ F_2^0(i) \cap \check{V}_3 &= F_3^0(\gamma(i)). \end{aligned} \tag{159}$$

For any $i, j \in I_2$ we have from formula (103) that

$$\begin{aligned} \mu(F_1(\varepsilon(i)) \mid F_1(\varepsilon(j))) &= \mu(F_1^0(\varepsilon(i)) \mid F_1^0(\varepsilon(j))) \\ \mu(F_3(\gamma(i)) \mid F_3(\gamma(j))) &= \mu(F_3^0(\gamma(i)) \mid F_3^0(\gamma(j))). \end{aligned} \tag{160}$$

We have the following statements.

Proposition 38 *Let $o \in I_2$. The following diagrams are commutative.*

1. *If E_1 is a cC_2 -space. Then*

$$\begin{array}{ccc} \mathcal{D}_{F_2(o)}(E_2) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(o)) \mid V_1) & \xrightarrow{\beta_*} & \mathcal{D}_{F_3(\gamma(o))}(E_3) \\ \mathbf{F} \otimes \mu(F_1(\varepsilon(o)) \mid V_1) \downarrow & & \downarrow \mathbf{F} \\ \mathcal{D}_{F_2^0(o)}(\check{E}_2) \otimes_{\mathbf{C}} \mu(F_1^0(\varepsilon(o)) \mid \{0\}) & \xrightarrow{(\check{\beta})^*} & \mathcal{D}_{F_3^0(\gamma(o))}(\check{E}_3). \end{array}$$

2. *If E_3 is a dC_2 -space. Then*

$$\begin{array}{ccc} \mathcal{D}_{F_2(o)}(E_2) \otimes_{\mathbf{C}} \mu(F_3(\gamma(o)) \mid \{0\}) & \xrightarrow{\alpha^*} & \mathcal{D}_{F_1(\varepsilon(o))}(E_1) \\ \mathbf{F} \otimes \mu(F_3(\gamma(o)) \mid \{0\}) \downarrow & & \downarrow \mathbf{F} \\ \mathcal{D}_{F_2^0(o)}(\check{E}_2) \otimes_{\mathbf{C}} \mu(F_3^0(\gamma(o)) \mid \check{V}_3) & \xrightarrow{(\check{\alpha})^*} & \mathcal{D}_{F_1^0(\varepsilon(o))}(\check{E}_1). \end{array}$$

3. *If E_1 is a cC_2 -space. Then*

$$\begin{array}{ccc} \mathcal{D}'_{F_3(\gamma(o))}(E_3) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(o)) \mid V_1) & \xrightarrow{\beta^*} & \mathcal{D}'_{F_2(o)}(E_2) \\ \mathbf{F} \otimes \mu(F_1(\varepsilon(o)) \mid V_1) \downarrow & & \downarrow \mathbf{F} \\ \mathcal{D}'_{F_3^0(\gamma(o))}(\check{E}_3) \otimes_{\mathbf{C}} \mu(F_1^0(\varepsilon(o)) \mid \{0\}) & \xrightarrow{(\check{\beta})^*} & \mathcal{D}'_{F_2^0(o)}(\check{E}_2). \end{array}$$

4. If E_3 is a dC_2 -space. Then

$$\begin{array}{ccc}
\mathcal{D}'_{F_1(\varepsilon(o))}(E_1) \otimes_{\mathbf{C}} \mu(F_3(\gamma(o)) \mid \{0\}) & \xrightarrow{\alpha_*} & \mathcal{D}'_{F_2(o)}(E_2) \\
\mathbf{F} \otimes \mu(F_3(\gamma(o)) \mid \{0\}) \downarrow & & \downarrow \mathbf{F} \\
\mathcal{D}'_{F_1^0(\varepsilon(o))}(\check{E}_1) \otimes_{\mathbf{C}} \mu(F_3^0(\gamma(o)) \mid \check{V}_3) & \xrightarrow{(\check{\alpha})^*} & \mathcal{D}'_{F_2^0(o)}(\check{E}_2).
\end{array}$$

5. If E_1 is a cfC_2 -space. Then

$$\begin{array}{ccc}
\mathcal{D}_{F_3(\gamma(o))}(E_3) & \xrightarrow{\beta^*} & \mathcal{D}_{F_2(o)}(E_2) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F} \\
\mathcal{D}_{F_3^0(\gamma(o))}(\check{E}_3) & \xrightarrow{(\check{\beta})^*} & \mathcal{D}_{F_2^0(o)}(\check{E}_2).
\end{array}$$

6. If E_3 is a dfC_2 -space. Then

$$\begin{array}{ccc}
\mathcal{D}_{F_1(\varepsilon(o))}(E_1) & \xrightarrow{\alpha_*} & \mathcal{D}_{F_2(o)}(E_2) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F} \\
\mathcal{D}_{F_1^0(\varepsilon(o))}(\check{E}_1) & \xrightarrow{(\check{\alpha})^*} & \mathcal{D}_{F_2^0(o)}(\check{E}_2).
\end{array}$$

7. If E_1 is a cfC_2 -space. Then

$$\begin{array}{ccc}
\mathcal{D}'_{F_2(o)}(E_2) & \xrightarrow{\beta_*} & \mathcal{D}'_{F_3(\gamma(o))}(E_3) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F} \\
\mathcal{D}'_{F_2^0(o)}(\check{E}_2) & \xrightarrow{(\check{\beta})^*} & \mathcal{D}'_{F_3^0(\gamma(o))}(\check{E}_3).
\end{array}$$

8. If E_3 is a dfC_2 -space. Then

$$\begin{array}{ccc}
\mathcal{D}'_{F_2(o)}(E_2) & \xrightarrow{\alpha^*} & \mathcal{D}'_{F_1(\varepsilon(o))}(E_1) \\
\mathbf{F} \downarrow & & \downarrow \mathbf{F} \\
\mathcal{D}'_{F_2^0(o)}(\check{E}_2) & \xrightarrow{(\check{\alpha})^*} & \mathcal{D}'_{F_1^0(\varepsilon(o))}(\check{E}_1).
\end{array}$$

Proof follows from formulas (159) and (160), constructions of two-dimensional direct and inverse images from section 5.6, two-dimensional Fourier transform, and corresponding properties of direct and inverse images and of Fourier transform on C_1 -spaces from section 4.8.

5.9 Two-dimensional Poisson formulas

In this section we obtain two-dimensional Poisson formulas: the Poisson formula I (subsection 5.9.1) and the Poisson formula II (subsection 5.9.2).

5.9.1

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of complete C_2 -spaces over a finite field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. By definition, there are order-preserving functions:

- (i) $\gamma : I_2 \longrightarrow I_3$ such that $\beta(F_2(i)) = F_3(\gamma(i))$ for any $i \in I_2$;
- (ii) $\varepsilon : I_2 \longrightarrow I_1$ such that $F_2(i) \cap V_1 = F_1(\varepsilon(i))$ for any $i \in I_2$.

Let $o \in I_2$. If we suppose that E_1 is a cC_2 -space, then from any $\mu \in \mu(F_1(\varepsilon(o)) \mid V_1)$ we canonically construct $\mathbf{1}_\mu \in \mathcal{D}'_{F_1(\varepsilon(o))}(E_1)$ in the following way. By definition, we have

$$\mathcal{D}'_{F_1(\varepsilon(o))}(E_1) = \varinjlim_{k \in I_1} \mathcal{D}'(V_1/F_1(k)) \otimes_{\mathbf{C}} \mu(F_1(\varepsilon(o)) \mid k). \quad (161)$$

By definition, $\mu(F_1(\varepsilon(o)) \mid V_1) \subset \mathcal{D}'(V_1/F_1(\varepsilon(o)))$. Therefore $\mu \in \mathcal{D}'(V_1/F_1(\varepsilon(o)))$. In formula (161) we take $k = \varepsilon(o)$. Then $\mu(F_1(\varepsilon(o)) \mid k) = \mathbf{C}$ and we put $1 \in \mu(F_1(\varepsilon(o)) \mid \varepsilon(o))$. Therefore μ gives $\mathbf{1}_\mu \in \mathcal{D}'_{F_1(\varepsilon(o))}(E_1)$ by formula (161).

Let $o \in I_2$. If we suppose that E_3 is a dC_2 -space, then from any $\nu \in \mu(F_3(\gamma(o)) \mid \{0\})$ we canonically construct $\delta_\nu \in \mathcal{D}'_{F_3(\gamma(o))}(E_3)$ in the following way. By definition, we have

$$\mathcal{D}'_{F_3(\gamma(o))}(E_3) = \varinjlim_{k \in I_2} \mathcal{D}'(F_3(k)) \otimes_{\mathbf{C}} \mu(F_3(\gamma(o)) \mid \{0\}). \quad (162)$$

In formula (162) we take $k = \gamma(o)$. Then we put $\delta_0 \in \mathcal{D}'(F_3(\gamma(o)))$, where $\delta_0(f) = f(0)$, $f \in \mathcal{D}(F_3(\gamma(o)))$. We put $\nu \in \mu(F_3(\gamma(o)) \mid \{0\})$ in formula (162). Thus we have constructed $\delta_\nu \in \mathcal{D}'_{F_3(\gamma(o))}(E_3)$.

We note that

$$\mathbf{F}(\mathbf{1}_\mu) = \delta_\mu \quad \text{and} \quad \mathbf{F}(\delta_\nu) = \mathbf{1}_\nu \quad (163)$$

under the maps

$$\mathbf{F} : \mathcal{D}'_{F_1(\varepsilon(o))}(E_1) \longrightarrow \mathcal{D}'_{F_1^0(\varepsilon(o))}(\check{E}_1) \quad \text{and}$$

$$\mathbf{F} : \mathcal{D}'_{F_3(\gamma(o))}(E_3) \longrightarrow \mathcal{D}'_{F_3^0(\gamma(o))}(\check{E}_3).$$

(We used that

$$\mu(F_1(\varepsilon(o)) \mid V_1) = \mu(F_1^0(\varepsilon(o)) \mid \{0\}) \quad \text{and} \quad \mu(F_3(\gamma(o)) \mid \{0\}) = \mu(F_3^0(\gamma(o)) \mid \check{V}_3).$$

Now we suppose that simultaneously E_1 is a cC_2 -space and E_3 is a dC_2 -space. Let $o \in I_2$. Then by any $\mu \in \mu(F_1(\varepsilon(o)) \mid V_1)$, $\nu \in \mu(F_3(\gamma(o)) \mid \{0\})$ it is well-defined *the characteristic function*

$$\delta_{E_1, \mu \otimes \nu} \stackrel{\text{def}}{=} \alpha_*(\mathbf{1}_\mu \otimes \nu) \in \mathcal{D}'_{F_2(o)}(E_2). \quad (164)$$

We have the following statement.

Lemma 6

$$\delta_{E_1, \mu \otimes \nu} = \beta^*(\delta_\nu \otimes \mu).$$

Proof follows from constructions and corresponding statements for the following admissible triple of C_1 -spaces:

$$0 \longrightarrow V_1/F_1(\varepsilon(k_2)) \longrightarrow \frac{F_2(k_1)}{F_2(k_2)} \longrightarrow F_3(\gamma(k_1)) \longrightarrow 0,$$

where $k_1 \geq o \geq k_2 \in I_2$ and $F_1(\varepsilon(k_1)) = V_1$, $F_3(\gamma(k_2)) = \{0\}$.

Now we have the following first two-dimensional Poisson formula.

Theorem 2 (*Poisson formula I*) . Let E_1 be a cC_2 -space, Let E_3 be a dC_2 -space. Let $o \in I_2$. Then for any $\mu \in \mu(F_1(\varepsilon(o)) \mid V_1)$, $\nu \in \mu(F_3(\gamma(o)) \mid \{0\})$ we have

$$\mathbf{F}(\delta_{E_1, \mu \otimes \nu}) = \delta_{\check{E}_3, \nu \otimes \mu}.$$

Proof follows from lemma 6, from formulas (163), and from proposition 38 about connection of direct and inverse images with two-dimensional Fourier transform. The theorem is proved.

Remark 23 Let $E = (I, F, V)$ be a C_2 -space over a field k . We construct a C_2 -space \overline{E} , which dominates a C_2 -space E . We define $\overline{E} = (I, F, V)$, but for every $i \leq j \in I$ the C_1 -space \overline{E}_{ij} is the maximal C_1 -space which dominates the C_1 -space E_{ij} , where \overline{E}_{ij} and E_{ij} are C_1 -spaces with underlying k -vector space $F(j)/F(i)$. (To construct for a C_1 -space (K, G, W) its maximal dominating C_1 -space (K', G', W) we denote by $Gr(W)$ the set of k -vector subspaces $U \subset W$ such that there are $k \leq l \in K$ with the property $G(k) \subset U \subset G(l)$. It is clear that $Gr(W)$ is a partially ordered set which is ordered by embeddings of k -subspaces. Then we put $K' = Gr(W)$, and the function G' is an embedding of the corresponding k -subspace to W .)

If

$$0 \longrightarrow E_1 \longrightarrow E_2 \longrightarrow E_3 \longrightarrow 0$$

is an admissible triple of C_2 -spaces, then

$$0 \longrightarrow \overline{E}_1 \longrightarrow \overline{E}_2 \longrightarrow \overline{E}_3 \longrightarrow 0$$

is also an admissible triple of C_2 -spaces.

We consider any admissible triple of C_2 -spaces over a finite field \mathbb{F}_q

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0,$$

where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$, and E_1 is a cC_2 -space, E_3 is a dC_2 -space. Let $g \in \text{Aut}_{C_2}(E_2)$ such that for any $i \in I_2$ there is $j \in I_2$ with the property $gF_2(i) = F_2(j)$. Then we have the following admissible triple of C_2 -spaces

$$0 \longrightarrow g\overline{E_1} \xrightarrow{g(\alpha)} g\overline{E_2} \xrightarrow{g(\beta)} g\overline{E_3} \longrightarrow 0, \quad (165)$$

where the C_2 -space $\overline{E_i}$ dominates the C_2 -space E_i ($1 \leq i \leq 3$) (see remark 23), $g\overline{E_2} = \overline{E_2}$, $g\overline{E_3} = \overline{E_2}/g\overline{E_1}$.

From (170) we have canonically

$$\mu(F_2(o) \mid gF_2(o)) = \mu(F_2(o) \cap gV_1 \mid gF_1(\varepsilon(o))) \otimes_{\mathbf{C}} \mu(g(\beta)(F_2(o)) \mid gF_3(\gamma(o))).$$

Then for any $a \in \mu(F_2(0) \mid gF_2(0))$ we consider $a = b \otimes c$, where

$$b \in \mu(F_2(o) \cap gV_1 \mid gF_1(\varepsilon(o))), \quad c \in \mu(g(\beta)(F_2(o)) \mid gF_3(\gamma(o))).$$

Under conditions and notations of theorem 2 we have

$$g\mu \in \mu(gF_1(\varepsilon(o)) \mid gV_1), \quad g\nu \in \mu(gF_3(\gamma(o)) \mid \{0\}).$$

We have $\tilde{g} = (g, a) \in \widehat{\text{Aut}_{C_2}(E_2)}_{F_2(o)}$. Then

$$R'_g(\delta_{E_1, \mu \otimes \nu}) = \delta_{gE_1, (g\mu \otimes b) \otimes (g\nu \otimes c)} = \delta_{gE_1, g\mu \otimes g\nu \otimes a}.$$

Corollary 1 *Let $(g, a) \in \widehat{\text{Aut}_{C_2}(E_2)}_{F_2(o)}$ such that for any $i \in I_2$ there is $j \in I_2$ with the property $gF_2(i) = F_2(j)$. Then we have*

$$\mathbf{F}(\delta_{gE_1, g\mu \otimes g\nu \otimes a}) = \delta_{\tilde{g}^{-1}(\check{E}_3), \tilde{g}^{-1}(\nu) \otimes \tilde{g}^{-1}(\mu) \otimes a}.$$

Proof follows from theorem 2 (Poisson formula I) and section 5.5.5.

5.9.2

Let

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0$$

be an admissible triple of complete C_2 -spaces over a finite field \mathbb{F}_q , where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$. By definition, there are order-preserving functions:

- (i) $\gamma : I_2 \longrightarrow I_3$ such that $\beta(F_2(i)) = F_3(\gamma(i))$ for any $i \in I_2$;
- (ii) $\varepsilon : I_2 \longrightarrow I_1$ such that $F_2(i) \cap V_1 = F_1(\varepsilon(i))$ for any $i \in I_2$.

If E_1 is a cfC_2 -space, then using remark 12 we have

$$\mathcal{D}(E_1) = \varprojlim_{i \geq j \in I_1} \mathcal{D}(F_1(i)/F_1(j)).$$

$F_1(i)/F_1(j)$ is a compact C_1 -space (for any $i \geq j \in I_1$). Therefore $1 \in \mathcal{D}(F_1(i)/F_1(j))$. Taking the projective limits, we obtain that in this case the following element is well-defined

$$\mathbf{1} \in \mathcal{D}(E_1). \quad (166)$$

If E_3 is a dfC_2 -space, then using remark 12 we have

$$\mathcal{D}(E_3) = \lim_{\substack{\leftarrow \\ i \geq j \in I_3}} \mathcal{D}(F_3(i)/F_3(j)).$$

$F_3(i)/F_3(j)$ is a discrete C_1 -space (for any $i \geq j \in I_3$). Therefore $\delta_0 \in \mathcal{D}(F_3(i)/F_3(j))$, where $\delta_0(0) = 1$. Taking the projective limits, we obtain in this case a well-defined element

$$\delta_0 \in \mathcal{D}(E_3). \quad (167)$$

We note that

$$\mathbf{F}(\mathbf{1}) = \delta_0 \quad \text{and} \quad \mathbf{F}(\delta_0) = \mathbf{1} \quad (168)$$

under the maps

$$\begin{aligned} \mathbf{F} &: \mathcal{D}(E_1) \longrightarrow \mathcal{D}(\check{E}_1) \quad \text{and} \\ \mathbf{F} &: \mathcal{D}(E_3) \longrightarrow \mathcal{D}(\check{E}_3). \end{aligned}$$

Now we suppose that simultaneously E_1 is a cfC_2 -space and E_3 is a dfC_2 -space. Let $o \in I_2$. Then an element

$$\delta_{E_1} \stackrel{\text{def}}{=} \alpha_*(\mathbf{1}) \in \mathcal{D}_{F_2(o)}(E_2) \quad (169)$$

is well-defined.

We have the following statement.

Lemma 7

$$\delta_{E_1} = \beta^*(\delta_0).$$

Proof follows from constructions and corresponding statements for the following admissible triples of C_1 -spaces (for any $i \geq j \in I_2$)

$$0 \longrightarrow \frac{F_1(\varepsilon(i))}{F_1(\varepsilon(j))} \longrightarrow \frac{F_2(i)}{F_2(j)} \longrightarrow \frac{F_3(\gamma(i))}{F_3(\gamma(j))} \longrightarrow 0,$$

where $\frac{F_1(\varepsilon(i))}{F_1(\varepsilon(j))}$ is a compact C_1 -space and $\frac{F_3(\gamma(i))}{F_3(\gamma(j))}$ is a discrete C_1 -space.

Now we have the following second two-dimensional Poisson formula.

Theorem 3 (Poisson formula II) . Let E_1 be a cfC_2 -space, Let E_3 be a dfC_2 -space. Let $o \in I_2$. Then

$$\mathbf{F}(\delta_{E_1}) = \delta_{\check{E}_3}.$$

Proof follows from lemma 7, from formulas (168), and from proposition 38 about direct and inverse images and two-dimensional Fourier transform. The theorem is proved.

We consider any admissible triple of C_2 -spaces over a finite field \mathbb{F}_q

$$0 \longrightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \longrightarrow 0,$$

where $E_i = (I_i, F_i, V_i)$, $1 \leq i \leq 3$, and E_1 is a cfC_2 -space, E_3 is a dfC_2 -space. Let $g \in \text{Aut}_{C_2}(E_2)$ such that for any $i \in I_2$ there is $j \in I_2$ with the property $gF_2(i) = F_2(j)$. Then we have the following admissible triple of C_2 -spaces

$$0 \longrightarrow g\overline{E_1} \xrightarrow{g(\alpha)} g\overline{E_2} \xrightarrow{g(\beta)} g\overline{E_3} \longrightarrow 0, \quad (170)$$

where the C_2 -space $\overline{E_i}$ dominates the C_2 -space E_i ($1 \leq i \leq 3$) (see remark 23), $g\overline{E_2} = \overline{E_2}$, $g\overline{E_3} = \overline{E_2}/g\overline{E_1}$.

Corollary 2 *Let $g \in \text{Aut}_{C_2}(E_2)$ such that for any $i \in I_2$ there is $j \in I_2$ with the property $gF_2(i) = F_2(j)$. Then we have*

$$\mathbf{F}(\delta_{gE_1}) = \delta_{\check{g}^{-1}(\check{E}_3)}.$$

Proof follows from theorem 3 (Poisson formula II) and section 5.5.5.

References

- [1] Algebraic Number Theory, Ed. J. W. S. Cassels and A. Fröhlich, Academic Press, L.-N.-Y., 1967
- [2] Beilinson A.A., Residues and Adeles, Funct. Anal. Pril., 14 (1980), no. 1, 44-45; English transl. in Func. Anal. Appl., 14, no. 1 (1980), 34-35.
- [3] Bruhat F., Distributions sur un groupe localement compact et applications à l'étude des représentations des groupes φ -adiques, Bull. Soc. Math. France, 89 (1961), 43-75.
- [4] Fimmel T., Parshin A.N., An introduction to the higher adelic theory, preprint 1999.
- [5] Gaitsgory D., Kazhdan D., Representations of algebraic groups over a 2-dimensional local field, Geom. and Funct. Analysis, 14(2004), n 3, 535-574; ArXiv math. RT/0302.174.
- [6] Huber A., On the Parshin-Beilinson Adeles for Schemes, Abh. Math. Sem. Univ. Hamburg, 61 (1991), 249-273.
- [7] Kac V. G., Peterson D. H., Spin and wedge representations of infinite-dimensional Lie algebras and groups, Proc. Nat. Acad. Sci. USA, 78(1981), 3308-3312.
- [8] Kapranov M. M., Double affine Hecke algebras and 2-dimensional local fields, J. Amer. Math. Soc., 14(2001), n 1, 239-262; e-print arXiv math.AG/9812021.

- [9] Kapranov M.M., Semiinfinite symmetric powers, e-print arXiv: math.QA/0107089.
- [10] Lefschetz S., Algebraic topology, AMS Colloquium Publications 27, Amer. Math. Soc. New York, (1942).
- [11] Osipov D.V., Central extensions and reciprocity laws on algebraic surfaces, (Russian) Mat. Sb. 196:10 (2005), 111-136; translation in Sb. Math. 196:10 (2005), 1503-1527; see also arXiv: math.NT/0501155.
- [12] Osipov D.V., n -dimensional local fields and adeles on n -dimensional schemes, Surveys in Contemporary Mathematics, Edited by N. Young, Y. Choi; London Mathematical Society Lecture Note Series, No. 347 (2007), pp. 131-164, Cambridge University Press; e-print math.AG/0508205.
- [13] Osipov D., Adeles on n -dimensional schemes and categories C_n , International Journal of Mathematics, vol. 18, no. 3 (2007), 269-279; e-print arXiv: math.AG/0509189
- [14] Parshin A.N., On the arithmetic of two-dimensional schemes I, Repartitions and residues, Math. USSR Izv. Vol.10 (1976) No.4, 695-729 (Engl.)
- [15] Parshin A. N., Chern classes, adeles and L-functions, J. für reine und angew. Math., 341 (1983), 174-192.
- [16] Parshin A.N., Vector Bundles and Arithmetical Groups I., Proc. Steklov Math. Institute, 208 (1995), 212-233; e-print arXiv: alg-geom/9605001.
- [17] Parshin A. N., Lectures in the University Paris-Nord, 1995-96.
- [18] Parshin A. N., Higher-dimensional local fields and L-functions, Invitation to higher local fields (Münster, 1999), 199–213 (electronic), Geom. Topol. Monogr., 3, Geom. Topol. Publ., Coventry, 2000, see also ArXiv: math.AG/0012151.
- [19] Parshin A. N., Harmonic analysis on adelic spaces and local fields, Mathematisches Forschungsinstitut Oberwolfach, Report No. 43/2005, Arakelov Geometry, September 2005, pp. 2471-2474.
- [20] Weil A., Basic Number Theory, Springer-Verlag, Berlin, 1967.

D.V. Osipov
 Steklov Mathematical Institute RAS
E-mail: *d_osipov@mi.ras.ru*

A.N. Parshin
 Steklov Mathematical Institute RAS
E-mail: *parshin@mi.ras.ru*